

Quality Control Of Impurities And Comparison Of Pharmacokinetic Parameters Of Angiotensin Receptor Antagonists

Dobrina Tsvetkova, Danka Obreshkova, Stefka Ivanova

Department of Pharmaceutical Chemistry, Medical University-Sofia, Faculty of Pharmacy
2 Dunav Str., Sofia 1000, BULGARIA
e – mail: dobrinka30@abv.bg

Danka Obreshkova, Stefka Ivanova

Department of Pharmacognosy and Pharmaceutical Chemistry
Medical University-Plovdiv, Faculty of Pharmacy
15A Vasil Aprilov, Plovdiv 4002, BULGARIA

Vladimir Yankov

Berlin Chemie
BULGARIA

Peter Atanasov

Clinic of Internal Diseases UMHATEM “N. I. Pirogov”
Sofia, BULGARIA

Abstract—The major risk for mortality from chronic heart and kidney disease worldwide is an inadequate treatment of hypertension. Therapy of hypertension becomes successfully by the developed in recent years a new class compounds – sartans (angiotensin II-receptor antagonists), specifically blocking renin angiotensin aldosteron system.

Benefits of sartans in hypertensive patients include reduction in left ventricular hypertrophy, improvement of diastolic function, decrease of ventricular arrhythmias, reduction of microalbuminuria, improvement of renal function, cardioprotective effect in patients with heart failure.

The impurity profiling of active pharmaceutical ingredients is an important quality control parameter.

Impurities are analysed by high performance liquid chromatography, thin layer chromatography, capillary electrophoresis and spectrophotometry.

For estimation of impurities of sartans the most applied methods, due to highest selectivity and specificity in separation and detection, are: HPLC-MS, HPLCMS/MS, HPLC-ESI/MS and HPLC-TOF/MS.

The important pharmacokinetic data of sartans are:

- 1) suitability for oral administration as to have physico-chemical properties, determining good pharmacokinetic behavior
- 2) suitability to the criteria of the rule of Christopher A. Lipinski (Rule 5): a) $M_r < 500$; b) H – donors (NH, OH) < 5 ; c) H – acceptors (N, O) < 10 ; d) $\log P < 5$: $\log P = 4.9$ (Candesartan); $\log P = 3.58$ (Eprosartan); $\log P = 4.52$ (Irbesartan); $\log P = 4.68$ (Losartan); $\log P = 4.31$ (Olmesartan); $\log P = 4.66$ (Telmisartan); $\log P = 3.68$ (Valsartan)

3) high plasma protein binding, which provide to be obtained once daily

4) Telmisartan is with the highest oral bioavailability and with the longest half-life

5) the bioavailability of other sartans is: Candesartan (3-11 %); Tasosartan (3-7 %); Zolasartan (20 %); Enoltasartan (36-72 %).

Keywords—sartans; impurities; pharmacokinetic parameters; metabolism.

I. INTRODUCTION.

Arterial hypertension is a widespread disease with more than 1 billion cases worldwide. The major risk for mortality from chronic heart and kidney disease is an inadequate treatment of hypertension. Stages in the development of arterial hypertension are: 1) I: Phase A (prehypertensive); Phase B (transient); 2) II: defined as SBP ≥ 160 mm Hg or DBP ≥ 100 mm Hg: Phase A (unsustainable); Phase B resistant; 3) III: Phase A (compensated); Phase B (uncompensated). Antihypertensive drugs are: beta-blockers, calcium antagonists, angiotensin-converting enzyme inhibitors, vasodilators and diuretics [1].

I. Pharmacological applications of sartans.

Sartans are applied in a number of heart diseases [2, 3]:

- 1) heart failure [4]; 2) prevention of atrial fibrillation in heart failure [5]; 3) myocarditis [6]; 4) hypertension with vascular hypertrophy [7]; 5) hypertension with left ventricular dysfunction [8]; 6) acute coronary syndrome [9]; 7) protection of the vascular endothelium [10]; 8) platelet aggregation [11]; 9) prevention of stroke [12]; 10) cerebral ischemia [11, 13]; 11) prevention of dementia [14]; 12) Alzheimer – sartans reduce the development of disease by 40 % compared to other antihypertensive agents [15] due to prevention of beta-amyloid-induced cognitive impairment [16];

13) Parkinson [17]; 14) migraine (Candesartan) [18]; 15) headache [19]; 16) inflammation: an antiinflammatory effect [20]; 17) type 2 diabetes: a) prevention of diabetes [21]; b) symptomatic hypertension in type 2 diabetes [22]; 18) kidney: symptomatic hypertension: a) renoprotective effect in the absence of diabetes [23]; b) renoprotective effect in type 2 diabetes [24, 25]; c) diabetic nephropathy [26]; d) prevention of progression of renal failure [27]; e) glomerulonephritis [28]; g) impaired renal function and hyperuricemia [29]; h) improvement of renal perfusion with hypercholesterolemia [30]; 19) liver fibrosis [31, 32] and portal hypertension [32]; 20) diabetic retinopathy [33]; 21) colitis [34]; 22) prevention of prostate cancer [35]; 23) inactivation of plasminogen [36].

Due to the synergistic effect the combination therapy of sartans with other antihypertensive drugs is more effective than the respective monotherapies. Sartans are used in combination with:

- 1) beta-blockers in hypertension: Propranolol [37]
- 2) calcium channel blockers in hypertension: Amlodipine besylate: Olmesartan Medoxomil [38], Telmisartan [39], Valsartan [40, 41]; Felodipine [42]; Nilvadipine [43]
- 3) angiotensin-converting enzyme inhibitors [44]; a) hypertension: Benazepril: Valsartan [45]; b) chronic heart failure [46]: Candesartan [47, 48]; c) diabetic nephropathy [49]; d) proteinuria [50]
- 4) thiazide diuretic Hydrochlorothiazide in hypertension: Candesartan cilexetil [51], Eprosartan, [52], Irbesartan [53, 54], Losartan [55, 56], Olmesartan Medoxomil [57], Telmisartan [58], Valsartan [59, 60]
- 5) statins in hypertension [61]: Rosuvastatin: Olmesartan, Irbesartan, Telmisartan [61], Simvastatin: Valsartan [62, 63].

II. Impurities of sartans.

During all stages of the pharmaceutical manufacturing process: development, production, stability testing and regulatory expectation of pharmaceutical formulations, the impurity profiling of active pharmaceutical ingredients is an important quality control parameter [64]. The importance of analysis of potentially toxic impurities is in order to increase the safety of drug therapy, quality and efficacy of pharmaceuticals [65].

Active ingredient related impurities include stereochemistry and functional group of active drug.

Process related impurities include chemicals, reagents, catalysts, residual solvents, synthetic intermediate products [66].

Stability related impurities are result of degradation or transformation of active pharmaceutical ingredient. International Conference of Harmonization have specified various limits for impurities in drug substances [67] and in drug products [68] as they may influence bioavailability, safety and efficacy of drugs [66].

Drug impurity profiling include identification and quantitative determination of related substances in drug substances and in pharmaceutical formulations [69].

Impurities are analysed by liquid and thin layer chromatography, capillary electrophoresis and spectrophotometry [64].

HPLC coupled with mass spectroscopy (MS) is the most applied method due to higher selectivity and specificity in separation and detection. Different HPLC methods are applied: LC-MS, LCMS/MS, LC-ESI/MS and LC-TOF/MS [70].

On TABLE 1. are presented data for empirical structures, molecular mass and melting point of sartans.

SARTAN	Empirical structure	Molecular mass	Melting point
	Abitesartan	C ₂₆ H ₃₁ N ₅ O	461.556
Azilsartan	C ₃₀ H ₂₄ N ₄ O ₈	568.5	157-159
Candesartan	C ₂₄ H ₂₀ N ₆ O ₃	440.45	183-185
Candesartan Cilexetil	C ₃₃ H ₃₄ N ₆ O ₆	610.66	163
Elisartan	C ₂₇ H ₂₈ ClKN ₆ O ₅	591.1	188-190
Eprosartan	C ₂₃ H ₂₄ N ₂ O ₄ S	424.52	260-261
Eprosartan Mesylate	C ₂₃ H ₂₄ N ₂ O ₄ S. CH ₃ SO ₃ H	520.625	248-250
Embusartan	C ₂₅ H ₂₄ FN ₅ O ₃	461.488	250-252
Fimasartan	C ₂₇ H ₃₁ N ₇ OS	501.65	260-263
Irbesartan	C ₂₅ H ₂₈ N ₆ O	428.23	180-181
Losartan	C ₂₂ H ₂₃ ClN ₆ O	422.92	183.5-184.5
Losartan Potassium	C ₂₂ H ₂₂ ClKN ₆ O	461.01	183.5-184.5
Mifasartan	C ₃₀ H ₃₀ N ₆ O ₃ S	554.667	192-195
Olmesarta Medoxomil	C ₂₉ H ₃₀ N ₆ O ₆	558.585	175-180
Pomisartan	C ₃₁ H ₃₀ N ₄ O ₂	490.596	176-178
Pratosartan	C ₂₄ H ₂₅ N ₆ O	413.499	184-187
Ripisartan	C ₂₃ H ₂₂ N ₈ O	426.476	235-238
Tazosartan	C ₂₃ H ₂₁ N ₇ O	411.459	243-246
Telmisartan	C ₃₃ H ₃₀ N ₄ O ₂	514.617	261-263
Saprisartan Potassium	C ₂₅ H ₂₂ BrF ₃ N ₄ O ₄ SK	611.431	272-275
Valsartan	C ₂₄ H ₂₉ N ₅ O ₃	435.519	116-117
Zolasartan	C ₂₄ H ₂₀ BrClN ₆ O ₃	555.814	421.5

TABLE 1. Empirical structure, molecular mass and melting point of sartans.

Related substances of some sartans are given in the following tables: Candesartan (TABLE 2.), Losartan (TABLE 3.); equal related substances for Losartan, Candesartan and Olmesartan. (TABLE 4.), Telmisartan (TABLE. 5.); Olmesartan (TABLE 6.), Valsartan (Fig. 1).

TABLE 2. Related substances of Candesartan.







	
Candesartan Methyl Ester	Candesartan Ethyl Ester
	
Trityl Candesartan	Desethyl Candesartan
	
Candesartan Cilxetil HCl Methoxy Analog	Desethyl Candesartan Cilxetil

TABLE 3. Impurities of Losartan.









	
Losartan EP Impurity A	Losartan EP Impurity C
	
Losartan EP Impurity D	Losartan EP Impurity H
	
Losartan EP Impurity I	Losartan EP Impurity J
	
Losartan EP Impurity K Losartan Aldehyde	Losartan Carboxylic Acid

TABLE 4. Impurities of Losartan, Candesartan, Olmesartan.






	
5-(4'-(Bromomethyl)(1,1'-bi-phenyl)-2-yl)-1-trityl-1H-tetrazole	p-(o-1H-tetrazol-5-ylphenyl)benzyl alcohol
	
p-(o-1H-tetrazol-5-ylphenyl)toluene	Triphenylmethanol
	
Triphenylmethyl Methyl Ether	

TABLE 5. Impurities of Telmisartan.














	
Telmisartan Impurity A	Telmisartan Impurity C
	
Telmisartan Impurity E	Telmisartan Impurity F
	
Telmisartan acyl glucuronide	

TABLE 6. Related substances of Olmesartan.

 <p>Olmesartan Methyl Ester</p>	 <p>Olmesartan Ethyl Ester</p>
 <p>Olmesartan Dimer Ester</p>	 <p>Olmesartan Methyl Ether</p>
 <p>Olmesartan Medoxomil Dehydro Impurity</p>	 <p>Olmesartan Methyl Ketone</p>
 <p>Ethyl 4-(1-Hydroxy-1-methylethyl)-2-Propyl-imidazole-5-Carboxylate</p>	 <p>Olmesartan (N-(Triphenyl-methyl)-5-(4'-bromomethyl-biphenyl-2-yl)tetrazole)</p>




Fig. 1. Related substances of Valsartan.

Initially the impurities of Valsartan were investigated by Nie, as it is isolated (S)-N-valeryl (N-{{2'-(1-methyl-tetrazol-5-yl)biphenyl-4-yl}-methyl}-valine.

In US Pharmacopeia are included:
 (R)-N-valeryl-N-{{2'-(1H-tetrazole-5-yl)biphenyl-4-yl} methyl} valine;
 (S)-N-butyryl-N-{{2'-(1H-tetrazole-5-yl)biphenyl-4-yl}-methyl}valine;
 (S)-N-valeryl-N-{{2'-(1H-tetrazole-5yl)biphenyl-4-yl}-methyl}valine benzyl ester.

Imp III, Imp IV and Imp V are obtained in the synthesis of Valsartan of Fig. 2. Imp III is obtained in pyridine and tetrahydrofuran at -5-0 °C from condensation of Imp I with 5-phenylvaleroyl chloride, formed by 5-phenylvaleric acid, by reaction with thionyl chloride at room temperature.

Imp IV and Imp V are formed from Imp I in the same manner from 4-pentenoic acid and 5-chlorovaleric acid, respectively, and Imp V is obtained after alkaline hydrolysis of the condensation product with sodium hydroxide in an aqueous medium [67].




Fig 2. Impurities of Valsartan.

Imp II (Fig. 2) is formed at the last step of the synthesis of Valsartan in a reductive cleavage of the phenyl sulfide group in the presence of Raney nickel in aqueous sodium hydroxide solution. Imp III and IV may result from desulphurization or free radical processes. 5-Chlorovaleric acid, from which is synthesized Imp II. Unreacted 5-chlorovaleric acid and 5-phenylthiovaleric acid reacted with Imp I and the product after alkaline hydrolysis gives Imp V (Fig. 3.) [71].




Fig. 3. Valsartan Imp III and Imp IV, obtained from Imp I and Imp II.

III. Methods for analysis of impurities of sartans.

Sodium azide is the impurity as is the precursor in the synthesis of Irbesartan, Candesartan and Valsartan and is determined by gradient HPLC on Hydro RP column (250 x 4.6 mm x 4 μm), room temperature and UV-detection at λ = 205 nm [72]. In Irbesartan substance sodium azide is analysed by ion chromatography [73].

Azilsartan impurities are investigated by HPLC method, using Inertsil ODS-3 column (250 x 4.6 mm x 5 μm) in gradient mode with mobile phase: mixture of acetonitrile and potassium dihydrogen orthophosphate buffer [74]. Gradient reverse-phase HPLC is developed for the quantitative determination of related compounds of Azilsartan kamedoxomil on YMC-Pack C₁₈ column (150 mm x 4.6 mm x 3 μm) [75].

Related substances Desethyl Candesartan cilexetil, 1N-Ethyl Candesartan cilexetil; 2N-Ethyl Candesartan cilexetil, 1N-Ethyl Oxocandesartan cilexetil; 2N-Ethyl Oxocandesartan cilexetil are analysed by HPLC/MS [76].

Is reported that Candesartan impurity: 2-ethoxy-1-[[2'-(1-ethyl-1H-tetrazol-5-yl)biphenyl-4-yl]me-thyl]-1H-benzimidazole-7-carboxylic acid ethyl ester is determined by HPLC with mass-spectrometry detection with electrospray ionisation (MS/ESI) [77].

Ultra High-Pressure Liquid Chromatography (UPLC) method is proposed for determination of Candesartan cilexetil impurities in tablet formulation. The chromatographic separation is performed on Waters Acquity UPLC system and BEH Shield RP₁₈ column using gradient elution of mobile phase A: 0.01 M phosphate buffer adjusted pH 3.0 with ortho-phosphoric acid and mobile phase B: acetonitrile : water = 95 : 5 % v/v, UV-detection at λ = 254 nm for Candesartan Ethyl Ester and Desethyl Candesartan Cilexetil and at λ = 210 nm for Triethylalcohol and Triphenylmethyl Methyl Ether impurities [78].

For assay of impurities of Eprosartan mesylate is reported HPLC method at Phenomenex Gemini C₁₈ column (250 mm x 4.6 mm x 5.0 μm), gradient elution with mobile phase: A: 10 mM ammonium acetate buffer (pH to 3.0) : acetic acid; B: acetonitrile, UV-detection [79].

Reversed-phase liquid chromatography method for the simultaneous determination of Eprosartan mesylate and its six impurities is applied. Separation is achieved in less than 7 min using a fused-core C₁₈ column (100 mm x 2.1 mm x 2.6 μm), gradient elution with mobile phase: 10 mM ammonium formate (pH = 3.0) : acetonitrile [80].

Eprosartan and related substances 4,4'-(5,5'-(1E,1'E)-3,3'-(4,4'-methylene-bis (thiophene-4,2-diyl))-bis-(2-carboxyprop-1-ene-3,1-diyl)-bis-(2-butyl-1H-imidazole-5,1-diyl)bis-(methylene) are estimated by HPLC/MS [81].

Stability indicating UPLC method for simultaneous determination of Eprosartan mesylate,

Hydrochlorothiazide and their impurities in tablets is developed by performing of chromatographic separation on Acquity HSS C₁₈ column (150 x 2.1 mm x 1.7 μm), gradient elution using acetonitrile and 10 mM disodium hydrogen phosphate buffer (pH = 5.5, adjusted with phosphoric acid), flow rate: 0.3 ml/min., UV-detection at λ = 274 nm [82].

Isocratic reverse phase high performance liquid chromatographic (RP-HPLC) method is used for the determination of 2-cyano-4'-bromomethyl biphenyl and 2-n-butyl-1,3-diazaspiro[4,4]-non-1-ene-4-one impurities of Irbesartan substance and in pharmaceutical dosage forms [83].

In RP-HPLC method for the estimation of process related impurities of Irbesartan the separation is carried out on Hypersil Octadecylsilyl C₁₈ column, (4.6 mm x 150 mm, 3 μm), column temperature: 25°C, mobile phase A: 0.55 % v/v orthophosphoric acid, pH = 3.2 with triethylamine; mobile phase B: solvent A : acetonitrile = 5 : 95 v/v, flow rate: 1.2 ml/min., UV-detection at λ = 220 nm. The gradient program is : time (min.) / % mobile phase B = 0/40, 10/40, 22/50, 26/50, 28/40 and 35/40 [84].

HPLC is developed for Losartan related substances by using of Kromasil 100-5C₁₈ column (250 x 4.6 mm x and 5 μm particle size), column temperature: 35 °C, mobile phase: 0.1 % phosphoric acid in water : acetonitrile = 50 : 50 v/v, UV-detection at λ = 220 nm [85].

Gradient HPLC method is developed for simultaneous quantitative determination of Losartan potassium and its included in European pharmacopoeia 7.0 related impurities [86]:

B ([2'-(1H-tetrazol-5-yl)biphenyl-4-yl]methanol)

C ([2-butyl-5-chloro-1-[[2'-(1H-tetrazol-5-yl)biphenyl-4-yl]methyl]-1H-imidazol-4-yl]methanol)

D (2-butyl-4-chloro-1H-imidazole-5-carbaldehyde)

E (5-(4'-methylbiphenyl-2-yl)-1H-tetrazole)

F (5-[4'-[[2-butyl-4-chloro-5-[(1-methylethyl)oxy]methyl]-1H-imidazol-1-yl]methyl]biphenyl-2-yl)-1H-tetrazole)

I 5-[4'-[[2-butyl-4-chloro-5-[(triphenylmethyl)oxy]methyl]-1H-imidazol-1-yl]methyl]biphenyl-2-yl)-1H-tetrazole)

G (triphenylmethanol) [87].

The reported chromatographic system is: ACCHROM ODS-C₁₈ column (250 mm x 4.6 mm x 5 μm), column temperature: 35 °C, mobile phase: acetonitrile : 0.1 % phosphoric acid under a gradient elution, flow rate: 1.0 ml/min., UV-detection at λ = 220 nm [87].

HPLC method is described for the estimation of Telmisartan related impurities in tablets formulation by using X-Bridge C₁₈ column (150 mm x 4.6 mm x 3.5 μm), mobile phase: 25 mM potassium dihydrogen phosphate : 10 mM 1-hexanesulphonic acid [88].

For the assay of Telmisartan related substances in tablets formulation, a gradient reverse phase ultra-performance liquid chromatographic (RP-UPLC) method is developed by using Waters Acquity BEH C₁₈ column (100 mm x 2.1 mm x 1.7 μm) and UV-detection at λ = 290 nm [89].

HPLC method for simultaneous quantification of low level impurities of Telmisartan and Hydrochlorothiazide in tablet dosage forms is reported [90].

UPLC method has been developed for simultaneous determination of Telmisartan impurities and Chlorthalidone impurities in their formulations. Chromatographic separation is carried out on an Acquity BEH Shield-RP₁₈ column (100 x 2.1 mm x 1.7 μm), column temperature: 25°C, mobile phase A: pH = 4.5 buffer : acetonitrile = 90 : 10 v/v; mobile phase B: pH = 4.5 buffers : acetonitrile = 20 : 80 v/v, flow rate: 0.3 ml/min, UV-detection at λ = 290 nm, injection volume: 3 μl.

pH = 4.5 buffer is prepared using 0.025 M potassium dihydrogen phosphate, 0.0027 M 1-hexanesulphonic acid sodium salt and 1 ml of triethylamine in milli-Q water. The gradient program is reported as follows: time (min.) / % mobile phase B: 0/20, 2/30, 5/45, 8/55, 10/80, 14/80, 14.1/20 and 18/20 [91].

Gradient HPLC with mass spectrometry is used for the detection of 5 impurities of Valsartan: (S)-N-(1-carboxy-2-methylprop-1-yl)-N-[2'-(1H-tetrazol-5-yl)-biphenyl-4-ylmethyl]amine (S)-N-(1-carboxy-2-methylprop-1-yl)-N-(5-phenylthio)pentanoyl-N-[2'-(1H-tetrazol-5-yl)-biphenyl-4-ylmethyl]amine (S)-N-(1-carboxy-2-methylprop-1-yl)-N-(5-phenyl)pentanoyl-N-[2'-(1H-tetrazol-5-yl)-biphenyl-4-yl-methyl]amine (S)-N-(1-carboxy-2-methylprop-1-yl)-N-4-pentanoyl-N-[2'-(1H-tetrazol-5-yl)-biphenyl-4-ylmethyl]amine (S)-N-(1-carboxy-2-methylprop-1-yl)-N-(5-hydroxy)pentanoyl-N-[2'-(1H-tetrazol-5-yl)-biphenyl-4-ylmethyl]amine [92].

HPLC condition are: RP₁₈, (250 mm x 4.6 mm x 5 μm); mobile phase A: 0.01 M KH₂PO₄ : 0.005 M K₂HPO₄; mobile phase B: water : acetonitrile = 1:4 v/v; gradient elution: A/B: 0/50, 20/70, 30/70, 40/80, 50/50, 60/50, 0.8 ml/min., λ = 210 nm [92].

Capillary zone electrophoresis is applied for chiral purity of Valsartan in tablets, where R-enantiomer of Valsartan is an impurity. Separations is carried out in a 50 μm, 64/56 cm fused-silica capillary, 25 mM phosphate buffer (pH = 8), containing 10 mM

acetyl- β -cyclodextrin as a chiral selector, applied voltage: 30 kV and temperature: 30 °C [93].

IV. Pharmacokinetic parameters of sartans.

Data on the pharmacokinetic parameters of sartans are summarized on TABLE 7. [94]: Candesartan [95], Irbesartan [96, 97], Olmesartan medo-

Sartan	Biological half-life [h]	Protein binding [%]	Bioavailability [%]	Renal clearance [%]	Hepatic clearance [%]
Candesartan cilexetil	4-9	>99	15-42	60	40
Eprosartan	5-9	97-98	13	30	70
Irbesartan	11-15	90-95	70	1	99
Losartan	2	98.7	33	10	90
EXP 3174	6-9	99.8	-	50	50
Olmesartan	14-16	>99	29	40	60
Telmisartan	24	>99	42-58	1	9
Valsartan	6	95	25	30	70

xomil [98], Valsartan [99].

TABLE 7. Pharmacokinetic parameters of sartans.

There are the following features in the pharmacokinetic behavior of sartans:

1) suitability for oral administration as of physico-chemical properties, determining good pharmacokinetic behavior

2) suitability to the criteria of the rule of Christopher A. Lipinski (Rule 5): a) $M_r < 500$; b) H – donors (NH, OH) < 5 ; c) H – acceptors (N, O) < 10 ; d) $\log P < 5$: $\log P = 4.9$ (Candesartan); $\log P = 3.58$ (Eprosartan); $\log P = 4.52$ (Irbesartan); $\log P = 4.68$ (Losartan); $\log P = 4.31$ (Olmesartan); $\log P = 4.66$ (Telmisartan); $\log P = 3.68$ (Valsartan) [100].

3) high plasma protein binding, which provide to be obtained once daily

Telmisartan is with the most high oral bioavailability and with the longest half-life. The bioavailability of other sartans is: Tasosartan (3-7 %); Zolasartan (20 %); Enoltasartan (36-72%) [100].

V. Metabolism of sartans.

Drug metabolism (biotransformation) is the major mechanism for maintaining homeostasis in the body exposure to the effects of drugs, pesticides, environmental contaminants. In biotransformation drug molecules are converted into metabolites which are with large polarity (by the addition of ionizable groups), with high degree of ionization at physiological pH, reduced capacity for protein binding, high molecular weight and larger sized molecules [101].

Under the influence of specific enzymes: cytochrome P450; microsomal aminooxidase with mixed function; Z-uridilphosphoglucuronil transferase; glutathione-S-transferase is going the phase of functionalization (I phase, non-synthetic chemical changes: oxidation, reduction, hydrolysis and hydration) in which in molecules are introduced active

groups: -OH; -COOH; NH_2 ; SH, which leads to increased hydrophilicity [101].

Factors affecting the metabolism are:

I. Chemical: lipophilicity, molecular size, degree of ionization.

II. Biological: gender, body type, age, hormones, diseases:

a) chronic liver diseases: reduce the processes of conjugation

b) chronic renal insufficiency: decreases the levels of cytochrome P₄₅₀ and slows down the metabolic changes

c) infectious diseases: increase the level of endogenous interferon component that inhibits certain metabolic pathways

d) acute alcohol impairment: inhibition of enzyme activity by contacting of ethanol with cytochrome P 450

e) chronic alcohol damage: induction of enzyme activity in I and II biotransformation phase [64].

III. Genetic: fast and slow metabolizers (acetylators).

Metabolites are excreted or participate in the phase II (conjugation): detoxifying reactions: acetylation, methylation, conjugation of a drug molecule or its metabolites from the first phase to the highly polar, water-soluble natural endogenous substances: glucuronic acid (O-, N-, S-, C- glucuronide); glutathione; amino acids: glycine, taurine, glutamic acid [101].

75 % of the drug metabolism is under an influence of Cytochrome P450 enzymes. CYP3A4 is responsible for 50 % of drug metabolism [101]. CYP2C is the second expressed P450 subfamily in human liver and CYP2C9 is the most highly expressed isoform [102].

The differences observed in lipid solubility, absorption/distribution, plasma protein binding, bioavailability, biotransformation, plasma half-life, and systemic elimination influence the duration of action, and efficacy of sartans. On the basis of the daily mg dose, the antihypertensive potency of sartans follows the sequence: Candesartan cilexetil > Telmisartan = Losartan > Irbesartan = Valsartan > Eprosartan [103]. Sartans are metabolized in the liver by the microsomal liver enzymes Cytochrome P450 2C9 (CYP2C9) and P450 3A4 (CYP3A4), but not by CYP 1A2, 2A6, 2B6, 2C8, 2C9, 2C19, 2D6, 2E1, 3A5 and 4A11.

After oral administration approximately 14 % of Losartan dose is converted to the pharmacologically active metabolite: 5-carboxylic acid, designated as EXP 3174 [104]. Metabolite is long-acting (6 to 8 h) and is 10-40 times more potent in blocking AT₁ receptors than Losartan [105]. Both Losartan and EXP 3174 are more than 98 % bound to plasma proteins [106]. The major metabolic pathway for Losartan is by the Cytochrome P450 (CYP) 3A4, 2C9 and 2C10 isoenzyme [104]. The two-step oxidation of Losartan to EXP 3174 is catalyzed by CYP3A4 [107, 108] and CYP2C9 in human liver microsomes [108]. Hydroxymethyl group at C5 is oxidized to a carboxyl group of the active metabolite EXP 3174 (Fig. 4.) [107].




Fig. 4. Oxidation of Losartan to EXP3179 and EXP3174.

Candesartan cilexetil, Embusartan, Losartan, Olmesartan medoxomil and Tasosartan are ester prodrugs and after absorption through the gastrointestinal tract are hydrolysed by esterases to its respective active metabolite: Candesartan-7-carboxylic acid, BAY 10-6735, EXP 3174, Olmesartan and Enoltasartan. The metabolites are more active than the corresponding prodrug: Candesartan-7-carboxylic acid: 30-100 times more than Candesartan cilexetil. Candesartan slowly is biotransformed to a very small extent by oxidation by CYP2C9 [95].

Irbesartan is metabolized in the liver by oxidation of P450 CYP2C9 and less by CYP3A4 and subsequent glucuronidation to Irbesartan main metabolite glucuronide (6 %) [96].

Tasosartan is biotransformed by oxidation of one or both methyl groups. Experimental oxidation is carried out by: a) SeO_2 , dioxane/water to the aldehyde derivative (6), which is oxidated from b): benzeneseleninic acid/ H_2O_2 in THF/tetrahydrofuran to a carboxyl derivative (3). The major metabolite of Tasosartan is Enoltasartan, which is active [109].




Fig. 4. Oxidation of Tasosartan.




Fig. 5. Oxidation of Valsartan to valeryl-4-hydroxy-valsartan.

Valsartan is oxidized from CYP2C9 to non-active metabolite 4-hydroxyvaleryl-Valsartan (Fig. 5.) [42].

Olmesartan medoxomil is hydrolysed [110] by carboxymethylenebutenolidase in the liver to the active metabolite RNH-6270 FK9 [111].

Paraoxonase 1 as a major bioactivating hydrolase for olmesartan medoxomil in human blood circulation [112].

Pratosartan is biotransformed to: (S)-(-)-2-(1-hydroxypropyl)-3-[2'-(1H-tetrazol-5-yl)biphenyl-4-ylmethyl]-5,6,7,8-tetrahydro-3H-cycloheptimidazol-4-one; 8- and 5-hydroxylated metabolite.

Candesartan cilexetil and its metabolite in human plasma and urine are quantified on Spherisorb S3P (phenyl) column (100 mm x 4.6 mm x 3 μm), mobile phase A: phosphate buffer (pH = 2.8) : acetonitrile : water = 20 : 20 : 60 v/v; phase B: phosphate buffer (pH = 2.8) : acetonitrile : water = 20 : 60 : 20 v/v, flow-rate was 1.0 ml/min., $\lambda_{\text{excitation}} = 265$ nm, $\lambda_{\text{emission}} = 395$ nm. The acetonitrile gradient A/B is: 0-9 min.: 80 %/20 %; 9-14 min.: 68 %/32 %, 14-25 min.: 80 %/20 % [113].

Losartan and its metabolite EXP 3174 are determined in biological material by HPLC, after liquid-liquid extraction on column ULTREMEX CN and UV-detection at $\lambda = 245$ nm [114]. Other HPLC with UV-detection method include analysis in human plasma, urine and dialysate. For plasma gradient method with phenyl analytical column, mobile phase: 25 mM potassium phosphate : acetonitrile = 70 : 30 v/v and fluorescence detection is used. For urine and dialysate an isocratic mobile phase: 25 mM potassium phosphate : acetonitrile = 60 : 40 v/v (pH = 2.2) is applied [115].

After extraction of Losartan and EXP 3174 from human plasma with ether, HPLC separation is carried out on Diamonsil C_{18} column, mobile phase: 0.02 M sodium dihydrogen phosphate buffer (pH = 2.35 with phosphoric acid) : acetonitrile = 57 : 43 v/v, flow rate: 0.5 ml/min., fluorescence detection at $\lambda_{\text{excitation}} = 250$ nm and $\lambda_{\text{emission}} = 370$ nm [116]. Yan Other applied HPLC method is with U1tremex CN

column (250 mm x 4.6 mm x 5 μ m), column temperature: 35 °C, mobile phase: isophosphoric acid (pH = 2.3; 0.015 M) : acetonitrile = 75 : 25 v/v; flow rate: 1.25 ml min., 35 °C, fluorescence detection at λ excitation = 250 nm and λ emission = 370 nm [117].

HPLC/MS with electrospray ionization with pneumatically-assisted nebulization is reported for the simultaneous determination of Losartan and EXP-3174 in human plasma, after extraction under acidic conditions by solid-phase extraction on a sorbent of copolymer, poly(divinylbenzene-co-N-vinylpyrrolidone). The analytes are separated on RP C₁₈ Phenomenex Synergy 4 POLAR-RP 80A column (50 mm x 2 mm), column temperature: t = 40 °C, isocratic mobile phase: 0.1 % triethylamine/0.1 % acetic acid (pH = 7.1) : acetonitrile = 65 : 35 v/v, flow rate: 0.7 ml/min. [118].

Losartan and Losartan acid in human plasma are estimated by LC-MS/MS method, after extraction from human plasma by solid phase extraction technique using Oasis HLB cartridge and following separation on Zorbax SB C₁₈ column [119].

In human plasma isocratic LC-MS-MS is applied by mobile phase: 0.1 % triethylamine: 0.1% acetic acid (pH = 7.1) : acetonitrile = 65 : 35 v/v [120].

HPLC-MS/MS for simultaneous assay of Losartan, EXP-3174 and Hydrochlorothiazide in human plasma is developed after liquid-liquid extraction, elution from C₁₈ column and detection with an API 3000 triple-quadrupole mass spectrometer using negative electrospray ionization and multiple reaction monitoring [121].

UPLC/MS/MS with multiple reaction monitoring in the negative ionization mode is described for Losartan, EXP-3174 and Hydrochlorothiazide in human plasma, after solid-phase extraction on Oasis HLB and analysis on BEH C₁₈ column, mobile phase: 1 % formic acid : acetonitrile = 15 : 85 v/v [122].

Losartan and its active metabolite EXP 3174 in human plasma and urine with limited plasma sample after liquid-liquid extraction are separated by RP HPLC/MS [123].

For analysis of Valsartan and its metabolite in human plasma is proposed solid phase extraction, followed by HPLC separation on RP C₁₈ Atlantis C18 column (100 mm x 3.9 mm), column temperature was kept to 40 °C, mobile phase: acetonitrile : phosphate buffer (5 mM, pH = 2.5) in gradient mode at flow rate: 1.30 ml/min., fluorescence detector at λ emission = 234 and λ excitation = 378 nm UV-detection at λ = 254 nm. Gradient elution mode is applied by the initial percentage of acetonitrile 32 % with a stepness of 4.5 %/min. to reach the 50 % [124].

CONCLUSION

The most applied methods for analysis of impurities of sartans, due to highest selectivity and specificity in separation and detection, are: HPLC-MS, HPLC/MS/MS, HPLC-ESI/MS and HPLC-TOF/MS.

The important pharmacokinetic data of sartans are: suitability to the criteria of the rule of Christopher Lipinski: LogP = 4.9 (Candesartan); LogP = 3.58 (Eprosartan); LogP = 4.52 (Irbesartan); LogP = 4.68 (Losartan); LogP = 4.31 (Olmesartan); LogP = 4.66 (Telmisartan); LogP = 3.68 (Valsartan) and high plasma protein binding.

Telmisartan is with the most high oral bioavailability and with the longest half-life.

Sartans are metabolized in the liver by the microsomal liver enzymes CYP2C9 and P450 3A4 (CYP3A4), but not by CYP 1A2, 2A6, 2B6, 2C8, 2C9, 2C19, 2D6, 2E1, 3A5 and 4A11.

REFERENCES

1. D.A. Taylor, and A.A. Abdel-Rahman, "Novel strategies and targets for the management of hypertension", *Adv. Pharmacol.*, vol. 57(1), pp. 291-345, November 2009.
2. S.G. Chrysant, "Angiotensin II receptor blockers in the treatment of the cardiovascular disease", *Clin. Ther.*, vol. 30(2), pp. 2181-2190, 2008.
3. M.C. Michel, C. Foster, H.R. Brunner, and L. Liu, "A systematic comparison of the properties of clinically used angiotensin II type 1 receptor antagonists", *Pharmacol. Rev.*, vol. 65(2), pp. 809-848, April 2013.
4. C.N. Gring, and G.S. Francis, "A hard look at angiotensin receptor blockers in heart failure", *J. Amer. Coll. Cardiol.*, vol. 44(9), pp. 1841-1846, November 2004.
5. J.S. Healey, A. Baranchuk, E. Crystal, C.A. Morillo, M. Garfinkle, S. Yusuf, and S.J. Connolly, "Prevention of atrial fibrillation with angiotensin-converting enzyme inhibitors and angiotensin receptor blockers: a meta-analysis", *J. Amer. Coll. Cardiol.*, vol. 45(11), pp. 1832-1839, June 2005.
6. X. Liu, X. Zhu, A. Wang, H. Fan, and H. Yuan, "Effects of angiotensin-II receptor blockers on experimental autoimmune myocarditis", *Int. J. Cardiol.*, vol. 137(3), pp. 282-288, November 2009.
7. R.D. Smith, H. Yokoyama, D.B. Averill, E.L. Schiffrin, and C.M. Ferrario, "Reversal of vascular hypertrophy in hypertensive patients through blockade of angiotensin II receptors", *J. Amer. Soc. Hypertens.*, vol. 2(3), pp. 165-172, May-June 2008.
8. R. Lakhdar, M.H. Al-Mallah, and D.E. Lanfear, "Safety and tolerability of angiotensin-converting enzyme inhibitor versus the combination of angiotensin-converting enzyme inhibitor and angiotensin receptor blocker in patients with left ventricular dysfunction: a systematic review and meta-analysis of randomized controlled trials", *J. Cardiac Failure*, vol. 14(3), pp. 181-188, April 2008.

9. S. Verma, M.M. Mamdani, M. Al-Omran, M. Melo, and J.L. Rouleau, "Angiotensin receptor blockers versus angiotensin converting enzyme inhibitors and acute coronary syndrome outcomes in elderly patients: a population-based cohort study (UMPIRE study results)", *J. Amer. Soc. Hypertens.*, vol. 1(4), pp. 286-294, July-August 2007.
10. A.U. Klingbeil, S. John, M.P. Schneider, J. Jacobi, R. Handrock, and R.E. Schmieder, "Effect of AT₁ receptor blockade on endothelial function in essential hypertension", *Amer. J. Hypertens.*, vol. 16(2), pp. 123-128, September 2003.
11. V. Papademetriou, "Inhibition of the renin-angiotensin-aldosterone system to prevent ischemic and atherothrombotic events", *Amer. Heart J.*, vol. 157(6) (Suppl. 1), pp. S24-S30, June 2009.
12. N. Tanahashi, "Stroke and renin-angiotensin system," *Nippon Rinsho*, vol. 64(6), pp. 1191-1199, June 2006.
13. N. Hosomi, A. Nishiyama, C.R. Ban, T. Naya, T. Takahashi, M. Kohno, and J. A. Koziol, "Angiotensin type 1 receptor blockage improves ischemic injury following transient focal cerebral ischemia", *Neurosci.*, vol. 134(1), pp. 225-231, 2005.
14. A. Fournier, R. Oprisiu-Fournier, J.M. Serot, O. Godefroy, J.M. Achard, S. Faure, H. Mazouz, M. Temmar, A. Albu, R. Bordet, O. Hanon, F. Gueyffier, J. Wang, S. Black, and N. Sato, "Prevention of dementia by antihypertensive drugs: how AT₁-receptor-blockers and dihydropyridines better prevent dementia in hypertensive patients than thiazides and ACE-inhibitors", *Exp. Rev. Neurother.*, vol. 9(9), pp. 1413-1431, September 2009.
15. P.R. Gard, and J.M. Rusted, "Angiotensin and Alzheimer's disease: therapeutic prospects", *Exp. Rev. Neurother.*, vol. 4(1), pp. 87-96, January 2004.
16. S. Takeda, N. Sato, D. Takeuchi, H. Kurinami, M. Shinohara, K. Niisato, M. Kano, T. Ogihara, H. Rakuji, and R. Morishita R, "Angiotensin receptor blocker prevented beta-amyloid-induced cognitive impairment associated with recovery of neurovascular coupling", *Hypertens.*, vol. 54(6), pp. 1345-1352, December 2009.
17. T.N. Grammatopoulos, S.M. Jones, F.A. Ahmadi, B.R. Hoover, L.D. Snell, J. Skoch, V.V. Jhaveri, A.M. Poczbott, J.A. Weyhenmeyer, and W.M. Zawada, "Angiotensin type 1 receptor antagonist Losartan, reduces MPTP-induced degeneration of dopaminergic neurons in substantia nigra", *Molecul. Neurodegener.*, vol. 2(1), pp. 1-17, January 2007.
18. E. Tronvik, L.J. Stovner, G. Helde, T. Sand, and G. Bovim, "Prophylactic treatment of migraine with an angiotensin II receptor blocker: a randomized controlled trial", *JAMA*, vol. 289(1), pp. 65-69, January 2003.
19. M. Etminan, M.A.H. Levine, G. Tomlinson, and P.A. Rochon, "Efficacy of angiotensin II receptor antagonists in preventing headache: a systematic overview and meta-analysis", *Amer. J. Med.*, vol. 112(8), pp. 642-646, June 2002.
20. L. Mascitelli, and F. Pezzetta, "Anti-inflammatory action of angiotensin-converting enzyme inhibitors and angiotensin receptor blockers", *J. Amer. Coll. Cardiol.*, vol. 47(4), pp. 889-891, February 2006.
21. J. Östergren, "Renin-angiotensin-system blockade in the prevention of diabetes", *Diabetes Res. Clin. Pract.*, vol. 76(Suppl. 1), pp. S13-S21, May 2007.
22. M.F.B. Braga, and L.A. Leiter, "Role of renin-angiotensin system blockade in patients with diabetes mellitus", *Amer. J. Cardiol.*, vol. 104(6), pp. 835-883, September 2009.
23. P.S. Leung, "Mechanisms of protective effects induced by blockade of the renin-angiotensin system: novel role of the pancreatic islet angiotensin-generating system in type 2 diabetes", *Diabetic Med.*, vol. 24(2), pp. 110-116, February 2007.
24. M. Arıcı, and Y. Erdem, "Dual blockade of the renin-angiotensin system for cardiorenal protection: an update", *Amer. J. Kidney Diseases.*, vol. 53(2), pp. 332-345, March 2009.
25. B.J. Hoogwerf, "Renin-angiotensin system blockade and cardiovascular and renal protection", *Amer. J. Cardiol.*, vol. 105(Suppl. 1), pp. 30A-35A, January 2010.
26. L. Lozano-Maneiro, and A. Puente-García. "Renin-angiotensin system blockade in diabetic nephropathy. Present evidences", *J. Clin. Med.*, vol. 4(11), pp. 1908-1937, November 2015.
27. M.Z. Molnar, K. Kalantar-Zadeh, E.H. Lott, J.L. Lu, S.M. Malakauskas, J.Z. Ma, D.L. Quarles, and C.P. Kovesdy, "Angiotensin-converting enzyme inhibitor, angiotensin receptor blocker use, and mortality in patients with chronic kidney disease", *J. Amer. Coll. Cardiol.*, vol. 63(7), pp. 650-658, February 2014.
28. F.P.C. Catapano, L. De Nicola, R. Minutolo, P. Zamboli, C. Gallo, and G. Conte, "Antiproteinuric response to dual blockade of the renin-angiotensin system in primary glomerulonephritis: meta-analysis and metaregression", *Amer. J. Kidney Diseases.*, vol. 52(3), pp. 475-485, September 2008.
29. A. Dang, Y. Zhang, G. Liu, G. Chen, W. Song, and B. Wang, "Effects of Losartan and Irbesartan on serum uric acid in hypertensive patients with hyperuricaemia in Chinese population", *J. Hum. Hypertens.*, vol. 20(1), pp. 45-50, January 2006.
30. A.R. Chade, M. Rodriguez-Porcel, S.J. Rippen-rop, A. Lerman, and L.O. Lerman, "Angiotensin II AT₁ receptor blockade improves renal perfusion in hypercholesterolemia", *Amer. J. Hypertens.*, vol. 16(2), pp. 111-115, February 2003.
31. Y. Nabeshima, S. Tazuma, K. Kanno, H. Hyogo, M. Iwai, M. Horiuchi, and K. Chayama, "Anti-fibrogenic function of angiotensin II type 2 receptor in CCl₄-induced liver fibrosis," *Biochem. Biophys. Res. Commun.*, vol. 346(3), pp. 658-666, August 2006.
32. U. Tox, and H.M. Steffen, "Impact of inhibitors of the renin-angiotensin-aldosterone system on liver fibrosis and portal hypertension", *Curr. Med. Chem.*, vol. 13(30), pp. 3649-3661, 2006.

33. A.K. Sjolie, "Prospects for angiotensin receptor blockers in diabetic retinopathy", *Diabetes Res. Clin. Pract.*, vol. 76(Suppl. 1), pp. S31-S39, May 2007.
34. O.I. Santiago, E. Rivera, L. Ferder, and C.B. Appleyard, "An angiotensin II receptor antagonist reduces inflammatory parameters in two models of colitis", *Regulatory Peptides*, vol. 146(1-3), pp. 250-259, February 2008.
35. H. Uemura, H. Ishiguro, and Y. Kubota, "Angiotensin II receptor blocker: possibility of antitumor agent for prostate cancer", *Mini Rev. Med. Chem.*, vol. 6(7), pp. 835-844, July 2006.
36. K.K. Koh, W.-J. Chung, J.Y. Ahn, S.H. Han, W.C. Kang, Y.-H. Seo, T.H. Ahn, I.S. Choi, and E.K. Shin, "Angiotensin II type 1 receptor blockers reduce tissue factor activity and plasminogen activator inhibitor type-1 antigen in hypertensive patients: a randomized, double-blind, placebo-controlled study", *Atherosclerosis*, vol. 177(1), pp. 155-160, November 2004.
37. J.H. Kim, J.M. Kim, Y.Z. Cho, J.H. Na, H. S. Kim, H.A. Kim, H.W. Kang, S.K. Baik, S.O. Kwon, S.H. Cha, Y.-J Kim, and M.Y. Kim, "Effects of Candesartan and Propranolol combination therapy versus Propranolol monotherapy in reducing portal hypertension", *Clin. Mol. Hepatol.*, vol. 20(4), pp. 376-383, December 2014.
38. S.G. Chrysant, M. Melino, S. Karki, J. Lee, and R. Heyrman, "The combination of Olmesartan medoxomil and Amlodipine besylate in controlling high blood pressure: COACH, a randomized, double-blind, placebo-controlled, 8-week factorial efficacy and safety study", *Clin. Ther.*, vol. 30(4), pp. 587-604, April 2008.
39. A. Sharma, A. Bagchi, S.B. Kinagi, Y.K. Sharma, V.P. Baliga, and C. Bollmall, "Results of a comparative, phase III, 12-week, multicenter, prospective, randomized, double-blind assessment of the efficacy and tolerability of a fixed-dose combination of Telmisartan and Amlodipine versus Amlodipine monotherapy in Indian adults with stage II hypertension", *Clin. Ther.*, vol. 29(12), pp. 2667-2676, December 2007.
40. T. Philipp, T.R. Smith, R. Glazer, M. Wernsing, J. Yen, J. Jin, H. Schneider, and R. Pospiech, "Two multicenter, 8-week, randomized, double-blind, placebo-controlled, parallel-group studies evaluating the efficacy and tolerability of Amlodipine and Valsartan in combination and as monotherapy in adult patients with mild to moderate essential hypertension", *Clin. Ther.*, vol. 29(4), pp. 563-580, April 2007.
41. D. Poldermans, R. Glazes, S. Kargiannis, M. Wernsing, J. Kaczor, Y.T. Chiang, J. Yen, R. Gamboa, and I. Fomina, "Tolerability and blood pressure-lowering efficacy of the combination of Amlodipine plus Valsartan compared with Lisinopril plus Hydrochlorothiazide in adult patients with stage 2 hypertension", *Clin. Ther.*, vol. 29(2), pp. 279-289, February 2007.
42. A. Nakashima, H. Kawashita, N. Masuda, C. Sauer, M. Niina, Y. Nagae, and K. Iwasaki, "Identification of cytochrome P450 forms involved in the 4-hydroxylation of Valsartan, a potent and specific angiotensin II receptor antagonist, in human liver microsomes", *Xenobiotica*, vol. 35(6), pp. 589-602, June 2005.
43. K. Noda, M. Ideishi, E. Tashiro, Y. Nakashima, M. Imamura, M. Seki, M. Fujino, T. Sou, M. Kohara, H. Kanaya, N. Saku, R. Kamei, M. Yamasaki, H. Sakai, N. Gondo, and K. Saku, "Efficacy and tolerability of Nilvadipine in combination with an angiotensin II receptor antagonist in patients with essential hypertension: a multicenter, open-label, uncontrolled study", *Curr. Ther. Res.*, vol. 64(3), pp. 151-166, March 2003.
44. F. Montecucco, and F. Mach, "Statins, ACE inhibitors and ARBs in cardiovascular disease", *Best Pract. Res. Clin. Endocrinol. Metabol.*, vol. 23(3), pp. 389-400, June 2009.
45. Y.S. Ke, Y.Y. Tao, H. Yang, G.H. Yu, "Effects of Valsartan with or without Benazepril on blood pressure, angiotensin II and endotoxin in patients with essential hypertension", *Acta Pharmacol. Sin.*, vol. 24(4), pp. 337-341, April 2003.
46. J.R. Kapoor, "Effect of combination angiotensin-converting enzyme and angiotensin receptor blocker therapy on heart failure mortality and morbidity", *Amer. J. Cardiol.*, vol. 101(5), pp. 744-745, March 2008.
47. M. White, S. Lepage, J. Lavoie, S. De Denus, M.-H. Leblanc, D. Gossard, L. Whittom, N. Racine, A. Ducharme, F. Dabouz, J.-L. Rouleau, and R. Touyz, "Effects of combined Candesartan and ACE inhibitors on BNP, markers of inflammation and oxidative stress, and glucose regulation in patients with symptomatic left failure", *J. Cardiac Failure*, vol. 13(2), pp. 86-94, March 2007.
48. J.P. Baguet, G. Barone-Rochette, and Y. Neuder, "Candesartan cilexetil in the treatment of chronic heart failure", *Vasc. Health Risk Manag.*, vol. 5(1), pp. 257-264, April 2009.
49. E.P. Havranek, "From black and white to shades of gray: race and renin-angiotensin system blockade", *J. Amer. Coll. Cardiol.*, vol. 51(19), pp. 1872-1873, May 2008.
50. M. MacKinnon, S. Shurraw, A. Akbari, G.A. Knoll, J. Jaffey, and H.D. Clark, "Combination therapy with an angiotensin receptor blocker and an ACE inhibitor in proteinuric renal disease: a systematic review of the efficacy and safety data", *Amer. J. Kidney Dis.*, vol. 48(1), pp. 8-20, July 2006.
51. K.P. Ohma, H. Milon, and K. Valnes, "Efficacy and tolerability of a combination tablet of Candesartan cilexetil and Hydrochlorothiazide in insufficiently controlled primary hypertension – comparison with a combination of Losartan and Hydrochlorothiazide", *Blood Press.*, vol. 9(4), pp. 214-220, 2000.
52. G.L. Plosker, "Eprosartan: a review of its use in hypertension", *Drugs*, vol. 69(17), pp. 2477-2499, 2009.

53. A.J. Lewin, and M.R. Weir, "Antihypertensive efficacy and tolerability of Irbesartan/Hydrochlorothiazide in hypertensive patients stratified by body mass index and type 2 diabetes mellitus status: a post hoc subgroup analysis of the Irbesartan/HCTZ blood pressure reductions in diverse patient populations trial", *Clin. Ther.*, vol. 30(12), pp. 2354-2365, December 2008.
54. C. Venkata, S. Ram, and T. Silfani, "A comparative analysis of the efficacy of Olmesartan medoxomil (O)/Hydrochlorothiazide (HCTZ), Valsartan (V)/HCTZ, Irbesartan (I)/HCTZ and Telmisartan (T)/HCTZ combinations", *Amer. J. Hypertens.*, vol. 17(5) (Suppl. 1), pp. S121, May 2004.
55. A.H. Gradman, W.E. Brady, L.P. Gazdick, P. Lyle, and R.K. Zeldin, "A multicenter, randomized, double-blind, placebo-controlled, 8-week trial of the efficacy and tolerability of once-daily Losartan 100 mg/Hydrochlorothiazide 25 mg and Losartan 50 mg/Hydrochlorothiazide 12.5 mg in the treatment of moderate-to-severe essential hypertension", *Clin. Ther.*, vol. 24(7), pp. 1049-1061, July 2002.
56. I. Lacourcière, J.M. Neutel, and H. Schumacher, "Comparison of fixed-dose combinations of Telmisartan/Hydrochlorothiazide 40/12.5 mg and 80/12.5 mg and a fixed-dose combination of Losartan/Hydrochlorothiazide 50/12.5 mg in mild to moderate essential hypertension: pooled analysis of two multicenter, prospective, randomized, open-label, blinded-end point (PROBE) trials", *Clin. Ther.*, vol. 27(11), pp. 1795-1805, November 2005.
57. S.G. Chrysant, M.A. Weber, A.C. Wang, and D.J. Hinman, "Evaluation of antihypertensive therapy with the combination of Olmesartan medoxomil and Hydrochlorothiazide", *Amer. J. Hypertens.*, 17(3), pp. 252-259, March 2004.
58. R. Fogari, A. Zoppi, A. Mugellini, P. Preti, M. Destro, A. Rinaldi, and G. Derosa, "Effectiveness of Hydrochlorothiazide in combination with Telmisartan and Olmesartan in adults with moderate hypertension not controlled with monotherapy: a prospective, randomized, open-label, blinded end point (PROBE), parallel-arm study", *Curr. Ther. Res.*, vol. 69(1), pp. 1-15, February 2008.
59. Y. Lacourcière, L. Poirier, D. Hebert, L. Assouline, P. Stolt, B. Rehel, and Y. Khder, "Antihypertensive efficacy and tolerability of two fixed-dose combinations of Valsartan and Hydrochlorothiazide compared with Valsartan monotherapy in patients with stage 2 or 3 systolic hypertension: an 8-week, randomized, double-blind, parallel-group trial", *Clin. Ther.*, vol. 27(7), pp.1013-1021, July 2005.
60. J.L. Pool, R. Glazer, N. Crikelair, and D. Levy, "The role of baseline blood pressure in guiding treatment choice: a secondary analysis of the use of Valsartan/Hydrochlorothiazide as initial therapy in hypertensive adults in a randomized, double-blind, placebo-controlled trial", *Clin. Drug Invest.*, vol. 29(12), pp. 791-802, 2009.
61. J.M. Lorenzen, H. Neunhöffer, S. David, J.T. Kielstein, H. Haller, and D. Fliser, "Angiotensin II receptor blocker and statins lower elevated levels of osteopontin in essential hypertension – results from the EUTOPIA trial", *Atherosclerosis*, vol. 209(1), pp. 184-188, March 2010.
62. S. Rajagopalan, F. Zannad, A. Radauceanu, R. Glazer, Y. Jia, M.F. Prescott, M. Kariisa, and B. Pitt, "Effects of Valsartan alone versus Valsartan/Simvastatin combination on ambulatory blood pressure, C-reactive protein, lipoproteins, and monocyte chemoattractant protein-1 in patients with hyperlipidemia and hypertension", *Amer. J. Cardiol.*, vol. 100(2), pp. 222-226, July 2007.
63. L.-C. Rump, E. Baranova, B. Okopien, M. Weisskopf, A. Kandra, and P. Ferber, "Coadministration of Valsartan 160 and 320 mg and Simvastatin 20 and 40 mg in patients with hypertension and hypercholesterolemia: a multicenter, 12-week, double-blind, double-dummy, parallel-group superiority study", *Clin. Ther.*, vol. 30(10), pp.1782-1793, October 2008.
64. P.D. Tzanavaras, "Recent advances in the analysis of organic impurities of active pharmaceutical ingredients and formulations: a review", *Curr. Org. Chem.*, vol. 14(19), pp. 2348-2364, November 2010.
65. A.C. Kogawa, and H.R.N. Salgado, "Impurities and forced degradation studies: a review", *Curr. Pharm. Anal.*, vol. 12(1), pp. 18-24, February 2016.
66. S.K. Nagpal, U. Ashutosh; R. Bhardwaj, T. Thakkar, "A review on need and importance of impurity profiling", *Curr. Pharm. Anal.*, vol. 7(1), pp. 62-70, February 2011.
67. International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use, ICH, Q3A: Impurities in New Drug Substances, 2006.
68. International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use, ICH, Q3B: Impurities in New Drug Products, October 2006.
69. D. Bartos, and S. Gorog, "Recent advances in the impurity profiling of drugs", *Curr. Pharm. Anal.*, vol. 4(4), pp. 215-230, November 2008.
70. V.K. Vyas, M. Ghate, R.D. Ukawala, "Recent advances in characterization of impurities – use of hyphenated LC-MS technique", *Curr. Pharm. Anal.*, vol. 6(4), pp. 299-306, November 2010.
71. A. Sampath, A.R. Reddy, B. Yakambaram, A. Thirupathi, M. Prabhakar, P.P. Reddy, and V. P. Reddy, "Identification and characterization of potential impurities of Valsartan, AT₁ receptor antagonist", *J. Pharm. Biomed. Anal.*, vol. 50(3), pp. 405-412, October 2009.
72. M. Gričar, and S. Andrenšek, "Determination of azide impurity in sartans using reversed-phase HPLC with UV-detection", *J. Pharm. Biomed. Anal.*, vol. 125(1), pp. 27-32, June 2016.
73. N.H. Subramanian, V.R.S. Babu, R.G. Jeevan, G. Radhakrishnan, "Matrix elimination ion chromatography method for trace level azide determination in Irbesartan drug", *J. Chromatogr. Sci.*, vol. 47(7), pp. 529-533, August 2009.

74. W. Zhou, Y. Zhou, L. Sun, Q. Zou, P. Wei, and P. Ouyang, "Structural elucidation of potential impurities in Azilsartan bulk drug by HPLC", *J. AOAC Int.*, vol. 97(6), pp.1552-1562, November-December 2014.
75. J. Sreenivasulu, P.V. Ramana, G.S.K. Reddy, Nagaraju CV, S.T. Rajan, S. Eswaraiyah, "A rapid novel HPLC method for estimation of eight related compounds in Azilsartan kamedoxomil and identification of degradation compounds by using LC-MS", *J. Chromatogr. Sci.*, vol. 53(9), pp. 1463-1474, October 2015.
76. A. Mohan, S. Shanmugavel, A. Goyal, B.R. Venkataraman, and D. Saravanan, "Identification, isolation and characterization of five potential degradation impurities in Candesartan cilexetil tablets", *J. Chromatogr.*, vol. 69(11), pp. 1211-1220, March 2009.
77. B. Raman, B.A. Sharma, G. Mahale, D. Singh, and A. Kumar, "Investigation and structural elucidation of a process related impurity in Candesartan cilexetil by LC/ESI-ITMS and NMR", *J. Pharm. Biomed. Anal.*, vol. 56(2), pp. 256-263, September 2011.
78. N.D.A. Kumar, K.S. Babu, U. Gosada, and N. Sharma, "A validated ultra high-pressure liquid chromatography method for separation of Candesartan cilexetil impurities and its degradants in drug product", *Pharm. Methods*, vol. 3(1), pp. 31-49, January 2012.
79. S. Cheekatla, R. Rupakula, B.N. Reddy, and M. Sharada, "Determination and characterization of process impurities for Eprosartan mesylate", *J. Pharm. Res.*, vol. 6(5), pp. 504-509, May 2013.
80. P.D. Kalariya, M.V.K. Talluri, V.D. Gaitonde, P.S. Devrukhakar, and R. Srinivas, "Quality by design: a systematic and rapid liquid chromatography and mass spectrometry method for Eprosartan mesylate and its related impurities using a superficially porous particle column", *J. Sep. Sci.*, vol. 37(16), pp. 2160-2171, August 2014.
81. C. Sun, J. Wu, D. Wang, and Y. Pan, "Characterization of a novel impurity in bulk drug eprosartan by ESI/MS(n) and NMR", *J. Pharm. Biomed. Anal.*, vol. 51(3), pp. 778-783, February 2010.
82. B. Sathesh, S. Pulluru, K. Nitin, and D. Saravanan, "Simultaneous determination of Eprosartan mesylate, Hydrochlorothiazide and their related compounds in pharmaceutical dosage forms by UPLC", *J. Liq. Chrom. Rel. Tech.*, vol. 34(17), pp. 1885-1900, October 2011.
83. P. Prabhu, and M. Muralidhar, "Development and validation of a high performance liquid chromatography method for simultaneous determination of Irbesartan and its related impurities in pharmaceutical tablets", *Int. J. Pharm. Sci. Drug Res.*, vol. 6(2), pp. 145-153, April-June 2014.
84. N. Goswami, "A validated stability-indicating liquid chromatographic method for determination of process related impurities and degradation behavior of Irbesartan in solid oral dosage", *J. Adv. Pharm. Technol. Res.*, vol. 5(1), pp. 33-40, January-March 2014.
85. M. Thapa, R. Nautiyal, M. Datar, and A.K. Singh, "Isolation, identification and characterization of process related impurities in Losartan potassium drug substance", *Asian J. Chem.*, vol. 22(6), pp. 4295-4298, 2010.
86. European pharmacopoeia, 7.0 version, General Monographs, Losartan potassium, 2009, 2382-2384.
87. S. Qiu, K. Liu, P. Ma, M. Wang, H. Chen, X. Xu, X. Hao, and Y. Wang. "Simultaneous analysis of Losartan potassium and its related impurities and degradation products in tablets using HPLC," *Curr. Pharm. Anal.*, vol. 11(1), pp. 25-34, 2015.
88. C.P. Kishore, V.B. Reddy, D.M. Kale, "Development and validation of stability indicating HPLC method for the estimation of Telmisartan related substances in tablets formulation", *Int. J. Res. Pharm. Sci.*, vol. 1(1), pp. 493-501, 2010.
89. V. Bhavani, T.S. Rao T, S.V.N. Raju, B. Madhusudan, and J. Begum, "Stability indicating UPLC method for the estimation of Telmisartan related substances in tablets formulation", *Int. J. Sci. Res. Publ.*, vol. 3(1), pp. 1-8, 2013.
90. M.R.M. Reddy, K.A. Praveen, R.V. Krishna, and H.S. Wasimul, "Stability-indicating HPLC method for simultaneous estimation of low level impurities of Telmisartan and Hydrochlorothiazide in tablet dosage forms, *Int. J. Pharm. Pharm. Sci.*, vol. 4(1), pp. 497-504, 2012.
91. B. Marineni, and T.S Reddy, "Simultaneous determination of Telmisartan impurities and Chlorthalidone impurities by UPLC". *Int. J. Pharm. Sci. Rev. Res.*, 26(2), pp. 226-230, May-June 2014.
92. A. Sampath, A.R. Reddy, B. Yakambaram, A. Thirupathi, M. Prabhakar, P. Pratap Reddy, V.P. Reddy, "Identification and characterization of potential impurities of Valsartan, AT₁ receptor antagonist", *J. Pharm. Biomed. Anal.*, vol. 50(3), pp. 405-412, 2009.
93. K.R. Lee, N.V.T. Nguyen, Y.J. Lee, S. Choi, J.S. Kang, W. Mar, K.H. Kim, "Determination of the R-enantiomer of Valsartan in pharmaceutical formulation by capillary electrophoresis", *Archiv. Pharm. Res.*, vol. 38(5), pp. 826-833, 2015.
94. J. Song, and M. White, "Pharmacologic, pharmacokinetic and therapeutic differences among angiotensin II receptor antagonists", *Pharmacother.*, vol. 20(2), pp. 130-139, February 2000.
95. J.Y. Jeon, Y.J. Im, Y. Kim, S.M. Han, M.J. Jo, D.H. Shin, J.S. Yoo, B.K. Moon, B.K. Kim, B.H. Lee, Y.H. Choi, B.S. Cho, H.Y. Jang, S.W. Chae, and M.G. Kim, "Pharmacokinetic properties and bioequivalence of Candesartan cilexetil in Korean healthy volunteers", *Drug Dev. Ind. Pharm.*, vol. 39(9), pp. 1296-1299, 2013.
96. M. Cánovas, F. Cabré, and F. Polonio, "Bioequivalence study of two oral formulations of Irbesartan 300 mg in healthy volunteers." *Drug Res. (Stuttg)*, vol. 64(1), pp. 53-56, January 2014.
97. S. Wittayalerpanya, P. Chariyavilaskul, N. Prompila, N. Sayankuldilok, and W. Eiamart, "Pharmacokinetics and bioequivalence study of Irbesartan tablets after a single oral dose of 300 mg in healthy Thai

- volunteers," *Int. J. Clin. Pharmacol. Ther.*, vol. 52(5), pp. 431-436, May 2014.
98. R. Kreutz, J. Bolbrinker, and M. Huber, "Pharmacokinetics of Olmesartan medoxomil plus Hydrochlorothiazide combination in healthy subjects", *Clin. Drug Investig.*, vol. 26(1), pp. 29-34, 2006.
99. M. Iqbal, A. Khuroo, L.S. Batolar, M. Tandon, T. Monif, and P.L. Sharma, "Pharmacokinetics and bioequivalence study of three oral formulations of Valsartan 160 mg: a single-dose, randomized, open-label, three-period crossover comparison in healthy Indian male volunteers", *Clin. Ther.*, vol. 32(3), pp. 588-596, March 2010.
100. K. Breithaupt-Grogler, C. Malerczyk, and G.G. Belz, "Pharmacodynamic and pharmacokinetic properties of an angiotensin II receptor antagonist-characterization by use of Schild regression technique in man", *Int. J. Clin. Pharmacol. Ther.*, vol. 35, pp. 434-441, 1997.
101. F.P. Guengerich, "Common and uncommon cytochrome P450 reactions related to metabolism and chemical toxicity". *Chem. Res. Toxicol.*, vol.14(6), pp. 611-650, June 2001.
102. S.A. Iwamura, T. Fukami, H. Hosomi, M. Nakajima, and T. Yokoi, "CYP2C9-mediated metabolic activation of Losartan detected by a highly sensitive cell-based screening assay", *Drug Metabol. Dispos.*, vol. 39(5), pp. 838-846, February 2011.
103. Z.H. Israili, "Clinical pharmacokinetics of angiotensin II (AT₁) receptor blockers in hypertension. *J. Hum. Hypertens.*, vol. 14(Suppl. 1), pp. S73-S86, April 2000.
104. D.A. Sica, T.W. Gehr, and S. Ghosh, "Clinical pharmacokinetics of Losartan", *Clin. Pharmacokinet.*, vol. 44(8), pp. 797-814, August 2005.
105. T. Tamaki, A. Nishiyama, S. Kimura, Y. Aki, M. Yoshizumi, H. Houchi, K. Morita, and Y. Abe, "EXP 3174: the major active metabolite of Losartan", *Cardiovasc. Drug Rev.*, vol. 15(2), pp. 122-136, April 1997.
106. G.P. Rossi, "Losartan metabolite EXP3179: an AT₁-receptor-independent treatment strategy for patients with the metabolic syndrome?", *Hypertens.*, vol. 54(1), pp. 710-712, August 2009.
107. C.H. Yun, H.S. Lee, H. Lee, J.K. Rho, H.G. Jeong, and F.P. Guengerich, "Oxidation of the angiotensin II receptor antagonist losartan (DuP 753) in human liver microsomes. Role of cytochrome P450 3A(4) in formation of the active metabolite EXP 3174", *Drug Metab. Dispos.*, vol. 23(2), pp. 285-289, February 1995.
108. R. Steams, P. Chakravarty, R. Chen, S. Chiu, "Biotransformation of Losartan to its active carboxylic acid metabolite in human liver microsomes. Role of cytochrome P450 2C and 3A subfamily members", *Drug Metab. Dispos.*, vol. 23(2), pp. 207-215, February 1995.
109. M. Elokda, M. Hassan, G.S. Friedrichs, S.-Y. Chai, B.L. Harrison, J. Primeau, M. Chlenov, and David L. Crandall, "Novel human metabolites of the angiotensin-II antagonist Tasosartan and their pharmacological effects", *Bioorg. Med. Chem. Lett.*, vol. 12(15), pp. 1967-1971, August 2002.
110. S.F. Ma, M. Anraku, Y. Iwao, K. Yamasaki, U. Kragh-Hansen, N. Yamaotsu, S. Hirono, T. Ikeda, and M. Otagiri, "Hydrolysis of angiotensin II receptor blocker prodrug Olmesartan medoxomil by human serum albumin and identification of its catalytic active sites", *Drug Metab. Dispos.*, vol. 33(12), pp.1911-1919, December 2015.
111. T. Ishizuka, I. Fujimori, M. Kato, C. Noji-Sakikawa, M. Saito, Y. Yoshigae, K. Kubota, A. Kurihara, T. Izumi, T. Ikeda, O. Okazaki, "Human carboxymethylenbutenolidase as a bioactivating hydrolase of Olmesartan medoxomil in liver and intestine. *J. Biol. Chem.*, vol. 285(16), pp.11892-11902, April 2010.
112. T. Ishizuka, I. Fujimori, A. Nishida, H. Sakurai, Y. Yoshigae, K. Nakahara, A. Kurihara, T. Ikeda, and T. Izum, "Paraoxonase 1 as a major bioactivating hydrolase for olmesartan medoxomil in human blood circulation: molecular identification and contribution to plasma metabolism", *Drug Metab. Dispos.*, vol. 40(2), pp. 374-380, February 2012.
113. H. Stenhoff, P.O. Lagerström, C. Andersen, "Determination of Candesartan cilexetil, Candesartan and a metabolite in human plasma and urine by liquid chromatography and fluorometric detection", *J. Chromatogr. B Biomed. Sci. Appl.*, vol. 731(2), pp. 411-417, August 1999.
114. A. Soldner, H. Spahn-Langguth, and E. Mutschler, "HPLC assays to simultaneously determine the angiotensin-AT₁ antagonist Losartan as well as its main and active metabolite EXP 3174 in biological material of humans and rats", *J. Pharm. Biomed. Anal.*, vol. 16, pp. 863-873, 1998.
115. D. Farthing, D. Sica, I. Fakhry, A. Pedro, and T.W.B. Gehr, "Simple high-performance liquid chromatographic method for determination of Losartan and E 3174 metabolite in human plasma, urine and dialysate", *J. Chromatogr. B*, vol. 704(1-2), pp. 374-378, December 1997.
116. L. Yang, D. Xia, and T. Guo, "Simultaneous determination of Losartan and its major metabolite (E 3174) in human plasma by HPLC-fluorescence method", *Zhongguo Yaofang*, vol. 20(32), pp. 2509-2511, 2009.
117. M.A. Ritter, C.I. Furtek, and M.-W. Lo, "An improved method for the simultaneous determination of Losartan and its major metabolite, EXP 3174, in human plasma and urine by high-performance liquid chromatography with fluorescence detection", *J. Pharm. Biomed. Anal.*, vol. 15(7), pp. 1021-1029, April 1997.
118. T. Iwasa, T. Takano, K.-I. Hara, and T. Kamei, "Method for the simultaneous determination of Losartan and its major metabolite, EXP 3174, in human plasma by liquid chromatography-electrospray ionization tandem mass spectrometry", *J. Chromatogr. B: Biomed. Sci. Appl.*, vol. 734(2), pp. 325-330, November 1999.
119. H.J. Shah, M.L. Kundlik, N.K. Patel, and C.N. Patel, "Rapid determination of Losartan and Losartan acid in human plasma by multiplexed LC-MS/MS", *J.*

Separat. Sci., vol. 32(20), pp. 3388-3394, October 2009.

120. B. Prasaja, L. Sasongko, Y. Harahap, Hardiyanti, W. Lusthom, and M. Grigg, "Simultaneous quantification of Losartan and active metabolite in human plasma by liquid chromatography-tandem mass spectrometry using Irbesartan as internal standard", J. Pharm. Biomed. Anal., vol. 49(3), pp. 862-867, April 2009.

121. F. Kolocouri, Y. Dotsikas, C. Apostolou, C. Kousoulos, and Y.L. Loukas, "Simultaneous determination of Losartan, EXP 3174 and Hydrochlorothiazide in plasma via fully automated 96-well-format-based solid-phase extraction and liquid chromatography-negative electrospray tandem mass spectrometry", Anal. Bioanal. Chem., vol. 387(2), pp. 593-601, January 2007.

122. P.A. Shah, P. Sharma, J.V. Shah, M. Sanyal, and P.S. Shrivastav, "Simultaneous analysis of Losartan, its active metabolite and Hydrochlorothiazide in human plasma by a UPLC-MS/MS method", Turk. J. Chem., vol. 39, pp. 714-733, January 2015.

123. M. Polinko, K. Riffel, H. Song, and M.-W. Lo, "Simultaneous determination of Losartan and EXP 3174 in human plasma and urine utilizing liquid chromatography/tandem mass spectrometry", J. Pharm. Bio-med. Anal., vol. 33(1), pp. 73-84, September 2003.

124. G. Iriarte, N. Ferreirós, I. Ibarondo, R.M. Alonso, M.I. Maguregi, L. Gonzalez, and R.M. Jiménez, "Optimization via experimental design of an SPE-HPLC-UV-fluorescence method for the determination of Valsartan and its metabolite in human plasma samples", J. Sep. Sci., vol. 29(15), pp. 2265-2283, October 2006.