

Evaluation Of The Antifungal And Antioxidant Activity Of *Citrullus Colocynthis* Against The Dieback Of Olive Tree In South Tunisia

Amira Bouzoumita^{1*}, Anis chikhoun² ¹, Nizar chairaa¹, Khaled Belouchette¹, Khawla chiheder³ and Ferchichi Ali¹

1- Arid and Oases Cropping Laboratory, Arid Area Institute-Medenine 4119

2-Université de Constantine et Université A/Mira de Béjaia

3-Technological institute of science Medenine(ISBAM) ,4119

*Faculty of Sciences of Tunis El Manar, 2292 Tunisia

Mednine-Tunisia

Corresponding author:

Arid and Oases Cropping Laboratory, Arid Area Institute-Medenine 4119

1-Amira Bouzoumita, Email:bouzoumita.amira@gmail.com

Abstract—To evaluate antioxidant activity and quantify total content of polyphenols, lipid and protein in two organs of plant extract of the *Citrullus colocynthis* (*C. colocynthis*) fruits and seeds. An qualitative and quantitative analysis of methanolic extracts, prepared from *C. colocynthis* fruits and seeds was carried out using standard methods. Estimation of their antioxidant activity was determined by free radical scavenging activity assay using 2,2-diphenyl-1-picrylhydrazyl and ferric reducing power and antibacterial activity from this three compounds. against a pathogens from olive a *Fusarium solani* and *Fusarium oxysporum* cause a new dieback of branches and roots of olive trees was observed in several orchards in southern Tunisia. Preliminary phytochemical screening of the methanol extracts showed that the content of the compounds varies among the two organs of cucurbits with the presence in different concentration of polyphenols, lipid and protein. Polyphenols from the bitter apple seeds have power over important antioxidant with a rate equal to 0.128295 g EAA / 100g monitoring the activity of polyphenols from fruit with a rate roughly equal (EAA 0.997614 g / 100g) in trotre, there are antioxidant activity outcome of seed proteins and fruit with the lowest value (0.5568 g EAA / 100g). While the lipid extract from seed lipid (0.99 g EAA / 100g) shows strong compared to that of fruit (0.089 g EAA / 100g) and relative to the protein with a slightly lower rate than that of polyphenols. These phytoconstituents ensure an interesting antifungal and antibacterial activity with a maximum inhibition rate of lipid, protein and polyphenolic extracts from the seeds that reaches 95%.

Keywords—antioxydant activity, antifungal activity, polyphenolic extracts, lipid and protein, antibacterial, , *Citrullus Colocynthis*, dieback, olive

1. Introduction

Current olive pest and infectious diseases caused by fungi are still a major threat to agriculture despite of the huge progress in of plant protection industry and pesticides. The disease control and management strategies are difficult to implement and so expensive (Pinto and Morais, 2000), (Trapero and Blanco, 1999, USEPA, 1996), since the pathogen lives in the ground and the application of conventional fungicides did not show any effectiveness (Sinclair and Hudler, 1998). The treatments applied are fundamentally cultural (Stipes, 2000). A simple method to protect the plant against these pathogens is particularly needed in developing countries due to relative widespread drug resistant microorganisms. Therefore, new antimicrobial substances must be developed and novel strategies should be investigated (Clardy and Walsh, 2004). Current research ocused on natural plant products as an effective source, based on their ethnobotany uses (Verpoorte and al, 2005). *Citrullus colocynthis* belongs to the family of Cucurbitaceae. Members of this family are generally dioeciously herbs which may be prostrate or climbing. *Citrullus colocynthis* had been extensively used in medicine since ancient times. medicallancient times. Traditionally, its fruit was used for the treatment of microbial diseases, jaundice, inflammation, ulcer, diabetes and urinary diseases in Asian and African countries (Saganuwan, 2010) and as insecticidal agent Rahuman and al, 2008. The aqueous extract of *Citrullus colocynthis* possesses an antidiabetic effect and antioxidant activity (Gurudeeban and Ramanathan, 2010). The purpose of this study is to highlight the beneficial effect of using of natural substances extracted from *Citrullus colocynthis* as bioprotective agents against olive dieback. Liquid medium dilution and direct contact methods were applied on potentially toxin-producing fungal strains isolated and identified from the olive trees. In this respect, phytochemicals from fruits and seeds have been shown to possess significant antioxidant

capacities. The antioxidant properties of fruit and seeds are depending on their content of phenolic compounds (flavonoids mainly), proteins and lipids. Polyphenols from the bitter apple seeds have power over important antioxidant with a rate equal to 0.128295 g EAA / 100g monitoring the activity of polyphenols from fruit with a rate roughly equal (EAA 0.997614 g / 100g) in turn, there are antioxidant activity outcome of seed proteins and fruit with the lowest value (0.5568 g EAA / 100g). While the antioxidant from seed lipid (0.99 g EAA / 100g) shows strong compared to that of fruit (0.089 g EAA / 100g) and relative to the protein with a slightly lower rate than that of polyphenols. Thus they could be utilized to control the quality of herbal preparations to ensure their antifungal and antimicrobial activities. Moreover, their effects of controlling of olive tree dieback of Plants caused by *Fusarium oxysporum* et *Fusarium solani*.

2. Materials and methods

2.1. Plant material

The fruits of *C. colocynthis* were collected at the stage of maturity, from September to November in a south location of Tunisia. Seeds were salvaged, dried, sheltered from whilst light and stored at ambient temperature till analysis. Five hundreds grams of each plant organ (fruit pulp and seeds) were powdered and passed through a 40-mesh sieve. The plant material was extracted with 95 % ethanol in a Soxhlet apparatus and ethanol was removed and dried at 40°C by a rotary evaporator.

2.2. Bacterial and fungal strains

2.2.1 Fungal support

PDA plates with the mycelium of the two strains of *Fusarium oxysporum* and *Fusarium solani* isolated and identified by microscopic and molecular techniques were incubated at 35 ° C for 4 to 5 days. In order to obtain pure cultures, each colony was subcultured repeatedly in new boxes of PDA and incubated at 35 ° C, in order to ensure growth and production of clean mycelium.

2.2.2. Microbial support

This support is composed of bacterial strains and reference strains as following: *Salmonella typhimurium* ATCC 1408 L2, *Enterococcus faecalis* A1 ATCC 29212 L3 A2, *Staphylococcus aureus* ATCC 25923 L6 A3 and *Escherichia coli* ATCC 35218 L5 A4.

2.3. Phytoscreening of *Citrullus Colocynthis*

2.3.1. Determination of total polyphenols by spectrophotometry

The concentration of total phenolics was measured by the method described by (Kim and al), with modifications. Briefly, an aliquot (1 ml) of standard solutions of gallic acid (20, 40, 60, 80 and 100 mg/ml) was added to a 25 ml volumetric flask containing 9 ml of deionized water (H₂O). After 5 min, 10 ml of 7%

Na₂CO₃ solution was added with mixing. One milliliter of Folin - Ciocalteu's phenol reagent, adjusted to 25ml with distilled water and was shaken thoroughly. After incubation for 90 min at 23 °C, absorbances of both blank and standard solutions were read at 750 nm. Total phenolic contents of samples were expressed as mg gallic acid equivalents (GAE)/100 g fresh weight. Absorbance of the pink color developed was determined at 765 nm. A calibration curve was prepared with quercetin and the results were expressed as mg quercetin equivalents /100g sample.

2.3.2. Radical-Scavenging Activity

DPPH radical-scavenging activity of sample extracts and standards (quercetin and BHT) were determined. The capacity of samples of *C. citrullus* seeds and fruit to scavenge the lipid-soluble 2, 2-diphenyl-1-picrylhydrazyl (DPPH) radical, which results in the bleaching of the purple color exhibited by the stable DPPH radical, is monitored at an absorbance of 515 nm. Radical scavenging activity of extracts and standards, which was interpolated from graph, constructed using percent inhibition against concentration of standards. An amount of 0.5 g of cold sample is added to 3 ml of methanol in a test tube, left in the dark and placed in an ice bath for one hour after making the samples. After one hour, the sample was taken from the tube with a Pasteur pipette, placed in 2 ml Eppendorf and centrifuged at a speed of 15,000 G, temperature of 4°C and for 10 min. The supernatant was recovered and put in two 5ml tubes. The DPPH was expressed as percentage inhibition by comparison with ascorbic acid standards (calibration curve constructed from 1.0 to 8.0 µg/mL).

2.3.3. Lipid extraction

Mature fruits were cut to get the seeds that have been finely ground and also conducted with crushed fruit extraction; using the mounting Soxhlet and hexane as solvent. After three hours, the recovered hexane is evaporated under reduced pressure at 50 ° C using a rotavapor [5], a viscous yellow oil having a foul odor was obtained.

2.3.4. Lipid assay

Total lipids were determined according to the method (Goldsworthy et al, 1972) using vanillin as reagent. The lipids form with hot sulfuric acid, together with the presence of vanillin and orthophosphoric acid, a pink complex. The assay is done on 100 µl aliquots of the lipid. The tubes are closed, shaken and placed for 10 min in a water bath. After heating to 100 ° C for 5 min, 200 µl of this mixture was added to 2.5 ml of sulphophosovanillin and vigorously stirred. After 30 min incubation in the dark, absorbances were read at 350 nm against a blank. The sulphophosovanillin is prepared as follows: dissolve 0.38 g of vanillin in 55 ml of distilled water and add 195 ml of orthophosphoric acid 85%. This reagent is stable for 3 weeks at 4 ° C in the dark. The lipid stock solution is freshly prepared from 2.5 ml edible oil.

2.3.5. Protein Extraction

From one gram of delipidated meal, proteins are extracted by a cold maceration for 24 hours

using 25 mL of three solvents (0.5 M NaCl, a toman solution pH = 7.4 and distilled water). After filtration, the volume of the extract is adjusted to 100 mL per each solvent.

2.3.6. Protein Assay

The protein assay was performed according to the (Bradford method 1976) in a 100 .ml liquot of 4 ml of Commassie Brilliant Blue (BBC; G 250; Merck) reagent is prepared in 50 ml of ethanol 95 °, then added by 100 ml of ortho phosphoric acid at 85% and made up to 1000 ml with distilled water .The shelf life of the reagent is 2 to 3 weeks at 4 ° C. The presence of protein in is revealed by the formation of blue dyes. Absorbance is read at 595 nm against a blank. Calibration curve is made from a bovine serum albumin solution titrating 1mg / ml.

2.4. Antifungal activity of phenolic, protein and lipid extracts of *colocynthis* in vitro

The antifungal activity of the studied extracts was assessed by different methods namely Well's method, disc method and direct contact method.

2.4.1. Wells' method (Double Layer)

In a Petri dish containing PDA medium, a thin layer of 0.6% PDA medium (3ml) containing 1ml of the spore suspension to be tested was spread on the surface. After solidification, 100 .ml of the sample is deposited into wells formed in the agar (Achemchem.F and al, 2004.6.2)

. Discs' method

9 mm diameter discs cut from Whatman No. 1, sterilized and impregnated with different concentrations of aqueous extracts of the tested biological materials (proteins, lipids and polyphenols) have been deposited gently on the surface of a previously inoculated medium. The inhibition diameters are measured around the disks after incubation in an incubator at 30-35 ° C for 24 to 48 h.

2.4.2. Direct contact method

Choice of solvent recovery polyphenolic extracts, lipid and protein ; DMSO seems to be the solvent that has no powerful antifungal potency over other solvent recovery (Yrjönen , 2004) , (Alvi, 2005), (Mohammedi , 2006,) and (Owagh and al ,2010). After incubation for 48 h, the culture medium (DMEM) of each cell was aspirated and replaced by 1 mL of a solution containing the following concentrations of DMSO (Dimehtylsulfoxide, DMSO; Mallinckrodt Baker Inc, Phillipsburg, NJ, USA): 0.05 mM (0.0004%); 0.1 mM (0.0008%); 0.3 mM (0.0024%); 0.5 mM (0.004%); or 1 mM (0.008%) mixed in plain DMEM. The control group was represented by DMSO-free DMEM. The cell were kept in contact with the solutions for 24 h and maintained in a humidified incubator at 37°C

under a 5% CO₂ and 95% air atmosphere during that period of time.

2.4.3. Antifungal test

The antifungal activity of the polyphenolic, lipid and protein extracts were carried out on two strains: (*Fusarium Oxysporum*, *Fusarium Solani*). They have been tested in vitro by direct contact method on PDA agar medium to determine the inhibition rate, comparing their actions to various concentrations of mycelial growth (Hussin and al, 2009), and a PDB liquid medium to determine the minimum inhibitory concentration (MIC), the fungicidal concentration and fungistatic concentration (FCS) (Derwich and al, 2010).

Determination of inhibition rate dilutions of *Citrullis Colocynthis* with a were prepared. To determine the inhibition rate, 3g of polyphenolic, lipid and protein extracts were introduced into a tube containing 10 ml of DMSO (300 mg / ml), 5 ml of the solubilized extract is then added to 05 ml of DMSO (150 mg / ml). The same procedure was applied for the preparation of concentrations from 18,75mg / ml to 300 mg / m . One ml of each extract is added to tubes containing 19ml of sterile liquid PDA medium. The mixture is homogenized at 45 ° C (Subrahmanyand al, 2001). Then, the mixture was put in Petri dishes of 90 mm (20ml / box) (Satish and al, 2010).

After solidification of the agar, the Petri dishes were divided into two parts, where a fungal strain was inoculated by a mycelial disk, taken from the youth culture of the fungus. The PDA without extract served as a control for each strain (Mishra and Dubey, 1994), (Khallil, 2001). The final concentrations of polyphenolic extracts used were calculated using the following equation

With : $C_f = C_i / 20$

C_f: final concentration of polyphenolic extracts, proteic and lipid in 1 ml of the PDA;

C_i = initial concentration of the polyphenolic extract, protein and lipid solubilized in DMSO [22]

Strains were incubated for 2 days to *Fusarium solani* and *Fusarium oxysporum* at a temperature of 30 ° C

The percentage inhibition of mycelial growth compared to the control, was calculated by the following formula: $PI(\%) = (A - B) / A \times 100$

With:

PI (%): inhibition rate expressed as percentage;

A: Diameter colonies in the "positive control" boxes;

B: Diameter of colonies in plates containing extract of polyphenols, lipids and proteins [23] Bajpai and al, 2010. The polyphenolic, lipid and protein extracts are considered very active when it has an inhibition between 75 and 100%, the fungal strain is considered

very sensitive when it has an inhibition between 50 and 74%, the fungal strain is called sensitive, moderately active when it has an inhibition between 25%, limited or not active when it has an inhibition between 0 and 24%, the fungal strain is called insensitive or resistant (Alcamo, 1984).

2.4.3.1. Liquid medium dilution method

Serial dilutions were made from the culture medium of the sample stock solution. To these dilutions was added a suspension of fungal microorganism. The preparation of these suspensions of fungal strains selected after sporulation of fungal strains (inoculated into 20 ml of PDA in Petri dishes after incubation at 30°C for 4 days), the spore cultures were recovered by adding 10 ml of sterile distilled water under vigorous stirring [(Solis-Pereira and al, 1993). A spectrophotometric assessment at 625nm (read at a wavelength of 625 nm) of the fungal suspension was carried out; to standardize the spore suspension to 106 spores / ml (Hussain and al, 2008).

2.5. Statistical Analysis

Data collected from all the experiments in this study were analyzed for statistical significance using analysis of variance (ANOVA). All samples were analyzed in three replications.

A. Means of treatments in each experiment were separated using Duncan's multiple range tests.

3. Results and discussion

3.1. Phenolic compounds, lipid and proteins contents of *Citrullus Colocynthis*

The seeds and fruits of *Citrullus* depict high contents of polyphenols, lipids and proteins, especially for the content of lipids assessed for the fruit

Figure 1: Content of phenolic compound from Coloquinte seeds and fruits

mg/ml Sample	Polyphenols	Proteins	Lipids
Seeds	36.63*±0,033	15.952*5±0,0083	51.666*5±0,018
Fruits	55.146*5±0,023	8.751 5±1,33	93.333*5±0,016

Table 1: Mean concentrations of extracted compounds

Values are the average of triplicate samples analysed individually. Star indicate significant differences.

Also, the contents are varying in seeds and fruits. Lipids are predominant compared to proteins and polyphenols in the two organs of *Citrullus*, fruits

showed the highest lipid contents. In another hand, the seeds and fruits have lowest protein contents. , however fruits exhibited lowest content compared to seeds with a threshold 0.00795 g/ml $P < 0.05$. The contents of polyphenols and proteins of *Colocynthis Citrullus* obtained in this study are similar to the results reported by (Pravinand al, 2013). However, lipid contents obtained in this study are different from those reported by (Simmons and al, 19(2006)

3.2. Determination of the antioxidant power of polyphenols

3.3. Antifungal tests with phenolic extracts, protein and lipid colocynthe

The results of the antifungal activity of phenolic compound are presented in (Figure 2).




Figure 2: percentage inhibition of polyphenols - fungus by the method well polyphenols extracts of seeds, : polyphenols extracts of fruits, FO : *Fusarium oxysporum*, *Fusarium solani*)

The analysis results showed that the antifungal activity is more important with the protein and lipid extracts from the seed of the fruit. The results of the antifungal activity of the lipid extract are presented in the following histogram (Figure 3)




Figure 3: Percent inhibition of lipid fungus by the method well

Polyphenolic extract from different parts including fruit a have been tested seeds on fungal strains (*Fusariumsolani*, *F.oxysporum*) after 5 days of incubation at 35 ° Cfor the antifungal activity, using the well method. Results obtained showed that the antifungal activity is higher with the phenolic fruit extract, followed by the extract of seeds and finally that of the antibiotic, showing moderately reduced mycelial growth.From the results shown in Figure4), there is a degradation of the pathogenic mycelium as it was depicted in Petri dishes




Figure 4: Inhibitory effect of polyphenol on *Fusarium solani*.

The results of the antifungal activity of the protein extract are presented in the following histogram(Figure 5)




Figure 5 : Percent inhibition of protein -fungi by the method wells.

The results Showed that the antifungal activity is higher with the seeds' protein extract than those of fruit and the antibiotic(Ampiciline), wherethe mycelial growth was reduced moderately.

the Minimum Inhibitory Concentrations (MICs) must be calculated in order to define the antifungal efficacy. (Tiwari et al, 2009) Results of calculations for polyphenolic , protein and lipid extracts results are shown in (Table 2).

Table 2: Minimum inhibitory concentrations (mg / ml) polyphenolic, lipid and protein extracts of *Colocynthis Citrullus* in liquid medium.

Fungal strains	MICs (mg/ml)		
	Lipids	Proteins	Polyphenols
<i>Fusariumoxysporum</i>	1.87	7.5	0.94
<i>Fusariumsolani</i>	0.94	3.75	0.94

The minimum inhibitory concentration (MIC), defined as the lowest concentration of polyphenols that has resulted in the complete inhibition of the growth of fungi [39] Jong and al, 2010, is ranging from 7.5 to 3.75 mg / ml for protein extract, from 1.87 to 0.94 mg / ml for the lipid extract and 0.94mg / ml for the polyphenol extract. *Solani* species strains have proven that fungal strains were more sensitive to the polyphenolic extract than the other two extracts.

3.4. Fungistatic and Fungicidal concentrations

The fungistatic (FCS) and fungicidal concentrations (FC) expressed in mg / ml are shown in (Table 3).

Table 3: Fungistatic (FCS) and fungicidal concentrations (FC) in mg / ml polyphenolic, lipid and protein extracts of *Citrullus Colocynthis*

Fungal strains	CF		CFS		CF		CFS	
	Lipids		Protein		Polyphenols			
<i>Fusariumoxysporum</i>	Ø	7.5	Ø	7.5	15	7.5		
<i>Fusariumsolani</i>	15	7.5	Ø	7.5	15	7.5		

Ø :empty set

Crops grown after obtaining there observed in the presence of polyphenolic, lipid and protein extracts of *Citrullus* on the strains tested. Protein extracts reveal fungistatic activity on fungal strains at a concentration of 7.5 mg / ml and no fungicidal activity while lipids exhibit fungicidal activity at a minimum concentration equal to 15mg / ml for the strain *Fusarium solani* only and fungistatic activity for the two fungals strains. As for the polyphenols, the fungicide concentration is 15 mg / ml and fungistatic concentration is 7.5 mg ml for the *Fusarium solani* strain. The studied strains do not have the same sensitivity. This could be due to the nature of the fungal strains and/or the wall, consisting of a complex network of proteins and polysaccharides and which varies in composition depending on the fungal species .Some polyphenolic compounds bind to microorganisms' proteins, blocking their enzymatic activities, (Mohamed Sham *et al.*, 2010) but also any DNA and RNA synthesis inhibition et d'ARN (Hadi, 2004).

3.5. DiskTest of antibacterial power

The results of the antibacterial activity in the following histogram (Figure 6)of the protein extracts using the disc method on agar medium showed that the proteins exhibit activity against four bacterial strains, namely. *Salmonella typhimurium* ATCC 1408 L2 A1, *Enterococcus feucalis* ATCC 29212 L3 A2, *Staphylococcus aureus* ATCC 25923 L6 A3 and *Escherichia coli* ATCC 35218 L5 A4.

A1 is more sensitive towards the protein extract. While there is no difference between the antibacterial activity of other bacterial strains. $P < 0.05$.




Figure 6: Percentage inhibition of the protein / bacteria Disk method (PR : protein, A : bacteria)




Figure 7: percentage inhibition of the lipid / bacteria by the method of discs (Gr : seeds, Fr : fruit, ATB : antibiotique, LIP : lipid)

From (Figure 7), the results of the antibacterial activity of the lipid extracts of *Citrullus* using the agar medium on disk method showed that lipids exhibit activity against four Bactrian strains. A1 bacterium is more sensitive lipids recovered from seeds than fruits. While there is a slight difference between the values of the antibacterial power $P < 0.05$. Moreover, the susceptibility of bacteria is almost equivalent to the antibiotic.




Figure 8: Percent inhibition of polyphenolic extracts versus bacteria by disc method

The results of the antibacterial activity of polyphenolic extracts using the PDA environment on disk method showed that polyphenols depicted antibacterial activity against 4 bacterial strains. A1 is

more sensitive to the polyphenolic extract from seeds than fruits. On the other hand, there is a slight difference between the antibacterial power for the bacterial strains while a low inhibition rate was observed for the antibiotic. From the previous data the polyphenol extracts from seeds showed a remarkable fungicidal effect against *Fusarium oxysporum* and *Fusarium solani* isolated from olive. In addition to that, these two strains do not show the same sensitivity, confirming the occurrence of non-phytotoxic and potentially effective compounds against pathogenic fungi (PRABAVATHY and al, 2006), (Chang and al, 2008).

This sensitivity is explained by the inhibition of fungal growth after transplanting discs in PDA media (direct contact method), which is similar to the findings on fungal strains of molds. The strong antifungal activity of the methanolic extract was also reported by (Hadizadeh and al, 2009) where the alcoholic extracts of *Citrullis Colocynthis* showed good antifungal activity (Figure 9).

According to (Yen Chang, 2008) In fact, the inhibition is due to the nature of the fungal strains of the wall composed of a complex network of proteins and polysaccharides and which varies in composition depending on the fungal species. (Domineco and al, 2005) reported that the disruption of this matrix by adsorption of phenolic compounds following the complexation polyphenols / membrane polysaccharides, may result in a reduction in the flow of internal and external layers of the wall which becomes defective and sensitive to osmotic lysis of antifungal agents. In addition, the antifungal activity of polyphenolic extracts could also be due to the fact that certain polyphenolic compounds bind to microorganisms proteins, blocking their enzymatic activities (Mohamed Sham and al, 2010). It may also add the inhibition of DNA synthesis and RNA [47] Hadi, 2004. In light of the results obtained, we can say that the lipid extracts of the seeds of *Citrullus colocynthis* exhibited slow mycelial growth. The work of (Amrouche A. and al, 2013) have shown that bitter apple oils are powerful natural antifungal and can be a very important source of constituents for plant protection used to eradicate fungal infections. The antimicrobial capacity of polyphenols act on the bacteria by the denaturation of cellular proteins (Hamdi and Ellouz, 1993) and alteration of the membranes. They may also inhibit the activity of symbiotic nitrogen-fixing bacteria by inhibiting the activity of digestive enzymes and / or precipitating nutritional proteins (Hättenschwiler and Vitousek, 2000).




Figure 9: Antifungal effect polyphenolic, lipid and protein extracts by the disk test

The antimicrobial capacity of polyphenols to inhibit the bacteria is explained by a fact of denaturation of cellular proteins (Hamdi and Ellouz, 1993) and alteration of the membranes (Ranalli, 1991). They can also inhibit the activity of symbiotic nitrogen-fixing bacteria by inhibiting the activity of digestive enzymes and / or precipitating nutritional proteins (Hättenschwiler and Vitousek, 2000).

Conclusion

Despite being a highly recalcitrant species, the olive tree is subject to a wide range of attacks of pathogens such as fungi, bacteria and viruses, which affect both the root, vegetative or reproductive system. However, fungal diseases in general and especially dieback cause the most significant damage often leading to the decline of the entire tree. The main causative agents of dieback disease are soil fungi, identification and purification of pathogens is the first step made in the context of our project is to classify each fungal strains and determine the range of the infecting the olive tree. The results of extraction and assessment of polyphenols, lipids and protein of *Citruscolocynthis* showed that their contents vary within the organs (fruit and seeds). It would be interesting to conduct further study on the various pathogenic fungal strains of the genus *Fusarium* responsible for certain deadly diseases of the olive tree and to appeal certain other biological products ensuring sustainable production of olive oils and very high quality and safety with a contamination level approaching zero tolerance. The purpose of studying the interaction of fungi and polyphenols, fats and proteins is to use these substances as biological agents in various cosmetics, pharmaceutical and drug alternative biopesticides against different pathogens of olive tree, as a part of a biological control.

References

- ABDEL HASSAN, I., ABDEL-BARRY, J.A. & MOHAMMEDA, S.T., 2000. The hypoglycaemic and antihyperglycaemic effect of *Citruscolocynthis* fruit aqueous extract in normal and alloxan diabetic rabbits. *Journal of Ethnopharmacology* 71, 325-330.
- AL-GHAITHI, F., EL-RIDI, M.R., ADEGHATE, E. & AMIRI, M.H., 2004. *Biochemical effects of Citruscolocynthis* in normal and diabetic rats. *Molecular and Cellular Biochemistry* 30, 1-7.
- Achemchem, F., J. M.-B. (2004). *purification et caractérisation d'une bactériocine anti-Listeria par Enterococcus faecium F-420 à partir du lait cru de chèvre*.
- Alcamo E-I. (1984). *Fundamentals of Microbiology*. Addison Wesley publishing company, . London , 310-341; 617-699.
- Ahmed Trigui, M. M. (2002). *Catalogue des variétés autochtones et types locaux*. 159p.
- Agri, M. (2005). *L'OLEICULTURE EN TUNISIE*
- Alcamo E-I. (1984). *Fundamentals of Microbiology*. Addison Wesley publishing company, . London , 310-341; 617-699.
- Ali RHOUMA, M. A. (2010). *First report of olive anthracnose caused by*. *Phytopathol. Mediterr* , 49, 95–98.
- Almi, D. (2010.). *Etude du pouvoir antioxydant des composés et extraits phénoliques issus des olives et sous produits de l'olivier*.
- Alavi S-H-R., Yassa N. et Fazeli M-R., 2005. *Chemical constituents and antibacterial activity of essential oil of Peucedanum ruthenicum M. Bieb. Fruits. SHR IJPS* 1 (4): 217-222
- A-R-M, K. (2001). . *Phytofungitoxic Properties in the aqueous extracts of some plants. Pakistan Journal of Biological Sciences* , 4 (4): 392-394.
- Balice V., C. O. (1984). *Acidic phenolic fraction of the juice of olives determined by gas chromatographie method. Grasas y Aceites*. . 25, 178 - 180.
- Boulila, M. (1994). *Le dépérissement de l'olivier en Tunisie : premiers résultats de recherche et perspectives d'avenir*.
- BENMEHDI, H., 2000. *Valorisation de certaines plantes médicinales à activité hypoglycémiantes comme la coloquinte. Mémoire de magistère en chimie organique appliquée. Université de Tlemcen. Algérie*
- BRUNETON, J., 1999. *Pharmacognosie, Phytochimie, plantes médicinales*. 3eme Ed, Tec & Doc Lavoisier, Paris, pp.1120.
- BENDJEDDOU, D., LALAOUI, K. & SATTA, D., 2003. *Immunostimulating activity of the hot water soluble polysaccharide extracts of Anacyclus pyrethrum, Alpinia galangal and Citruscolocynthis. Journal of Ethnopharmacol* 88, 155-160.
- BOUKEF, K. & SOUISSI, H.R., 1982. *Contribution à l'étude des plantes médicinales en médecine populaire en Tunisie. Rev. Soc. Pharm* 3, 34- 35.
- BNOUHAM, M., ZIYYAT, A., MEKHFI, H., TAHRI, A. & LEGSSYER, A., 2006. *Medicinal plants with potential antidiabetic activity - A review of ten*

years of herbal medicine research (1990-2000). *International Journal of Diabetes and Metabolism* 14, 1-25

- BENMEHDI, H., 2000. Valorisation de certaines plantes médicinales à activité hypoglycémiantes comme la coloquinte. Mémoire de magistère en chimie organique appliquée. Université de Tlemcen. Algérie.

- Bajpai V-K., S. S.-S. (2010). Antifungal activity of leaf essential oil and extracts of *Metasequoia glyptostroboides*. *Journal. American. Oil. Chemistry. Soc* , (87): 327-33

- Braga F-G., B. M.-L.-M.-L.-O.-O.-S. (2007). *Antileishmanial and antifungal activity of plants used in traditional medicine in Brazil*. *Journal of Ethnopharmacology* , 111: 396-402

- B., N. (2006). *Systématique. In : les champignons parasites des plantes cultivées : biologie, systématique, pathologie, maladies*. Centre de publication universitaire, Tunis , 119-195 .

- Bchard, J. (2009). Groupement CID/SCET-SCOM/ADI/CRC Sogema, Plan de Gestion des Pesticides et des Engrais et Guide de Bonnes pratiques.

- Bcherrawi, N. ((2002)). *Extraction des polyphénols des margines. Mémoire de fin d'études d'Ingénieur Agronome. Université Libanaise*. 111 pages.

- BENKADA, M. E. (2007). *Essai de lutte biologique in vitro par utilisation de Trichoderma Sp.A L'égard De Ascochyta Pinodella Agent De l'anthracnose du pois (Pisum Sativum L.)*.

- Benkada, M. Y. (2007). *Evaluation in vitro de l'effet antifongique de l'extrait acétonique et l'huile essentielle extraits de Thymus vulgaris L contre Fusarium oxysporum*.

- Bennett, A. W. (1992). The potential role of *Trichoderma viride* in the integrated control of *Botrytis fabae*. *The Mycologist* , 6:199-201.

- Bennett., K.-C. K. (1992). *Medical Mycology*. Lea & Febiger. Philadelphia and London.

- Bezert. (1996). *Action de Bacillus et d'actinomycètes sur les champignons du bleuissement de bois*.

- Boughalleb-M'Hamdi N*, S. M. (2011). *Screening of fungi implicated in the dieback of olive trees (Olea europea) in Chebika's area*. *Research in Plant Biology*, 1(4):33-38, 2011 .

- Boulila, M. (1994). *Le dépérissement de l'olivier en Tunisie : premiers résultats de recherche et perspectives d'avenir*.

- Boulila, M. M. (1994). *Inventaire des maladies de l'olivier en Tunisie*. *Bulletin OEPP*, 24 (4): 817-823.

- Boulila, M. (2001). *Maladies de l'olivier en Tunisie*. *Olivae* , 85:22-25.

- Braga F-G., B. M.-L.-M.-L.-O.-O.-S. (2007). *Antileishmanial and antifungal activity of plants used in traditional medicine in Brazil*. *Journal of Ethnopharmacology* , 111: 396-402.

- Borhade Pravin1*, Deshmukh Tushar1, Patil Vijay1 and Khandelwal Kishanchnad2, : *REVIEW ON Citrulluscolocynthis*, *IJRPC* 2013, 3(1). ISSN: 2231-2781

- Catalano, L. F. (1999). *Polyphenols in olive mill wastewaters and their depuration plant effluents: a comparison of the Folin-Ciocalteu and HPLC methods*. *Agrochimica*, , 43, pp. 193
- Chang C-W., C. W.-L.-T.-S. (2008). *Antibacterial activities of plant essential oils against Legionella pneumophila*. *Water Research* , 42: 78-286-205.

- CAMURATI F., A. L. (1984). *Le acque di vegetazione della lavorazione delle olive: tecnologie ed economie di recupero di sottoprodotti*. *Riv. Ital. Sostanze Grasse*, , 61, 283- 292.

- Catalano, L. F. (1999). *Polyphenols in olive mill wastewaters and their depuration plant effluents: a comparison of the Folin-Ciocalteu and HPLC methods*. *Agrochimica*, , 43, pp. 193-205.

- Chang C-W., C. W.-L.-T.-S. (2008). *Antibacterial activities of plant essential oils against Legionella pneumophila*. *Water Research* , 42: 78-286.

- Choi, G. J., Jang, K. S., Kim, J. S, Lee, S.W., Cho, J. Y., Cho, K. Y. and Kim, J. C. (2004), *In vivo antifungal activities of 57 plant Extracts against six plant pathogenic fungi*. *Plant Pathol. J.*, 3, 184-191

- CHANG, H., CHENG, Y., WU, C., CHANG, S., CHANG, T. & SU, T., 2008. *Antifungal activity of essential oil and its constituents from Calocedrusmacrolepisvar.fomiosana Florin leaf against plant pathogenic fungi*. *Bioresource Technology* 99, 6266-6270.

- Cichelli A., S. M.((1984). *composti fenolici delle olive e dell'olio di oliva*. *Riv. Merceol*.

- Cowan M-M. (1999.). *Plant Products as Antimicrobial Agents*. *Clinical Microbiology Reviews*. , 12 (4): 564-582.

- Dalile, A. Amrouche, H. Benmehdi, K. Fellah, H. A. Moussaoui, D. Chabane Sari., 2013 : *Essai in vitro de l'effet antifongique synergique des huiles extraites des graines de Citrulluscolocynthis L., Linumusitatissimum L. et de Nigellasativa L. sur Aspergillus ochraceustoxinogène* : *Revue des Régions Arides - Numéro Spécial - n° 35 (3/2014)*

- Dangles O, S. C. (1992). *Two very distinct types of anthocyanin complexation: Copigmentation and inclusion*. *Tetrahedron Lett.* , 33: 5227-30.

- Derwich E., B. Z. (2010). *GC/MS Analysis and antibacterial activity of the essential oil of Mentha pulegium grown in Morocco*. . Res. Journal of Agricultural and Biology , 6 (3):191-198.
- Djebali, N. (2008). *Etude des mécanismes de résistance de la plante modèle Medicago truncatula vis-à-vis de deux agents pathogènes majeurs des légumineuses cultivées :Phoma medicaginis et Aphanomyces euteiches*.
- DARWISH-SAYED, M., BALBAA, S.I. & AFIFI, M.S.A., 1973. Nitrogenous base of the different organs of Citrulluscolocynthis. *PlantaMedica*24(3), 260-265.
- Domenico T., Francesco C., Maria G-S., Vincenza V., Mariateresa C-D., Antonella S., Gabriela M. et Giuseppe B., 2005. Mechanisms of antibacterial action of three monoterpenes.*Antimicrobial Agents and Chemotherapy.*, 49 : 2474-2478.
- DUKE, J.A., 1978. The quest for tolerant germplasm. *ASA Special Symposium 32, Crop tolerance to suboptimal land conditions*. Am. Soc. Agron. Madison, WI: 1-61.
- Dulger B. et Gonuz A. 2004. Antimicrobial activity of some Turkish medicinal plants. *Pakistan journal of biological sciences.*, 7 (9) : 1559-1562
- Elaboration des etudes de faisabilite technique, et d'evaluation environnementale ainsi que la supervision technique de mise en œuvre des travaux etdu plan de gestion ENVIRONNEMENTALE DU PROJET ARBORICULTURE EN ZONES PLUVIALES (2009).
- FAO, O. d. (2013). *importance de l'oleiculture dans le monde*.
- FAO, O. d. (2012). *importance de l'oleiculture dans le monde*
- Fiorentino, A. G. ((2003)). *Environmental effects caused by olive mill wastewaters: toxicity comparaison of low-molecular-weight phenol components*. *Journal of Agricultural Food* .
- .
- Gamel, T. e. ((1999)). *Effect of methanol extracts of rosemary and olive vegetable water on the stability of olive oil and sunflower oil*. . *Grasas y Aceites*, , 50, pp. 345-350.
- GEBHARDT, R., 2003. *Antioxidative, antiproliferative and biochemical effects in Hep G2 cells of a homeopathic remedy and its constituent plant tinctures tested separately or in combination*. *Arzneimittelforschung*53, 823-830.
- GURUDEEBAN, S., RAJAMANICKAM, E., RAMANATHAN, T. & SATYAVANI,K., 2010. *Antimicrobial activity of citrulluscolocynthisin gulf of mannar*. *International Journal of Current Research* 2, 078-081.
- . G. Mazza& Y. S. Vefioglu .,1992. *Anthocyanins and other phenolic compounds in fruits of red-flesh apples* *Food Chemistry* 43 (1992) 113-117
- Hadi M., 2004. *La quercétine et ses dérivés: molécules à caractère pro- oxydant ou capteurs de radicaux libres; études et applications thérapeutiques*. Thèse doctorat. Université Louis Pasteur: 155 P.
- Hagerman AE, R. K. (1998). *High molecular weight plant polyphenolics (tannins) as biological antioxidants*. *Food Chem* , 46: 1887-92.
- Hamdi M., E. P. ((1993)). *Treatment of detoxified olive mill wastewater's by anaerobicfilter and aerobic fluized bed processes*. *Environ. Technol.* , 14, 183-188.
- HADIZADEH, F., EBRAHIMZADEH, M.A., HOSSEINZADEH, H., MOTAMEDSHARIATY,V., SALAMI, S. & BEKHRADNIA, A.R., 2009. *Antidepressant and antioxidant activities of some 2-benzoxazolinone derivatives as bupropion analogues*. *Pharmacologyonline* 1: 331-335
- Hattenschwiler S. et Vitousek P.M., 2000. *The role of polyphenols in terrestrial ecosystem nutrient cycling*. *Trends in Ecology and Evolution*. 15: 238-243
- Heinze, K. (2013). *Tunisie : des huiles d'olives bio de première qualité*.
- Hilal, A., & Chemseddine, M. H. (2002). *Lutte integree contre les ravageurs de l'olivier*.
- Hmouni A., H. M. ((1996)). *Résistance de Botrytis cinerea aux benzimidazoles et aux dicarboximides dans les cultures abritées de tomate en Tunisie*. *OEPP/EPPO Bull.* , 26, p:697–705.
- Hossain M-A., Z. I. (2008.). *Chemical composition and anti-fungal properties of the essential oils and crude extracts of Orthosiphon stamineus Benth*. *Industrial crops and products* , (27): 328–334.
- Huang D., O. B.-L. (2005). *The Chemistry behind Antioxidant Capacity Assays*. *Journal of Agricultural & Food Chemistry*. , 53: 1841-1856.
- Hussin N-M., M. R.-R.-A.-F.-A.-N.-K. (2009). *Antifungal activity of extracts and phenolic compounds from Barringtonia racemosa L. (Lecythidaceae)*. *African Journal of Biotechnology* , 8 (12): 2835-2845.
- JAMOUISSI, B. (1955). *Les maladies de déperissement des Agrumes*.
- Jardak T, T. M. (2007). *techniques de production en oléiculture-protection phytosanitaire-conseil oléicole international* p : 287-299.
- Jardak.J, Jerray.A, & Mahjoub.M. (2004). *La protection integree de l'oleiculture dans les pays de l'afrisie du nord*.
- Khallil A-R-M., 2001. *Phytofungitoxic Properties in the aqueous extracts of some plants*. *Pakistan Journal of Biological Sciences*. 4 (4): 392-394
- K.C. Highnam, G.J. Goldsworthy.,1972.Regenerated corpora cardiaca and hyperglycemic factor in Locusta migratoriaGeneral and Comparative

Endocrinology, Volume 18, Issue 1, February 1972, Pages :83-88

- KARRAY, B. (2002). *La filière huile d'olive en Tunisie: performances et stratégies*.
- KHLIF, N. G. (2001). *Caractérisation technologique des variétés d'olivier cultivées en tunisie*.
- l'Agriculture, M. d. (2006). *Direction Générale des Etudes et du Développement Agricole 2004-2005 Tunisie*.
- l'OLive, A. F., & AFIDOL. (2012). Le marché de l'huile d'olive :situation et perspective. *Alexandra PARIS – Responsable communication & économie* .
- Léger, C. (1999). Co-produits de l'huilerie d'olive: les composés phénoliques et leurs propriétés biologiques. . *OCL. Oléagineux, Corps gras, Lipides* , 6, pp : 60-63.
- l'olive, A. F. (2012).
- Lopez-Escudero FJ, D. R.-L. (2004). *Evaluation of olive cultivars for resistance to Verticillium dahliae. Europ. J. Plant. Pathol.*, 110: 79-85.
- Madi A. (2010.). Caractérisation et comparaison du contenu polyphénolique de deux plantes médicinales (Thym et Sauge) et la mise en évidence de leurs activités biologiques. Thèse magister. Université Mentouri Constantine:. 116 P
- MARZOUK, B., MARZOUK, Z., HALOUI, E., FENINA, N., BOURAOU, A. & AOUNI, M., 2010. Screening of analgesic and anti-inflammatory activities of *Citrullus colocynthis* from southern Tunisia. *J. Ethnopharmacol* 128, 15-19.
- M., B., M., M., & M.N, R. M. (1993). la maladie du pourridié-agaric observé dans des oliverais tunisiennes. *Bull OEPP/OPPE Bull.* .23:447-448.
- Mujica M-V., G. M. (2009). Importance of the extraction method in the quantification of total phenolic compounds in *Phaseolus vulgaris* L.Venzuela. . *Interciencia* , Vol. 34, N° 9: 650-654.
- Mohammedi Z., 2006. Etude du pouvoir antimicrobien et antioxydant des huiles essentielles et flavonoïdes de quelques plantes de la région de Tlemcen. *Thèse magistère*. Université de Tlemcen: 104 P
- M., H. (1991). Nouvelle conception d'un procédé de dépollution biologique des margines, effluents liquides de l'extraction de l'huile d'olive. Thèse de l'université de Provence. *Marseille ,France* , 166 p.
- Macheix J-J., F. A. (2005). Les composés phénoliques dans la plante : structure, biosynthèse, répartition et rôles. In: les polyphénols en agroalimentaire, Cheynier V., Sarni Manchado P. *Lavoisier, Paris*: , 510 P.
- Madi A. (2010.). Caractérisation et comparaison du contenu polyphénolique de deux plantes médicinales (Thym et Sauge) et la mise en évidence de leurs activités biologiques. Thèse magister. Université Mentouri Constantine:. 116 P.
- MAATOOQ, G.T., EL-SHARKAWY, S.H., AFIFI, M.S. & ROSAZZA, P.N., 1997.C-p-Hydroxybenzoylglyco-flavones from *Citrullus colocynthis*. *Phytochemistry* 44(1), 187.
- Mishra A-K. et Dubey N-K. (1994). . Evaluation of Some Essential Oils for Their Toxicity against Fungi Causing Deterioration of Stored Food Commodities. . *Applied and Microbiology.* , 60 (4): 1101-1105.
- Mohamed Sham S., Hansi H., Priscilla D. et Kavitha T., 2010. Antimicrobial activity and phytochemical analysis of selected Indian folk medicinal plants. *International Journal of Pharma Sciences and Research (IJPSR)* Vol. 1(10): 430-434.
- Mémoire de mastère en biologie GACEM Mohamed Amine :2011., Contribution à l'étude de l'activité antifongique et antimycotoxinogène des extraits méthanolique et aqueux des graines de *Citrullus colocynthis* sur la croissance de quelque moisissure d'altération de blé tendre stocké. FACULTE DES SCIENCES DE LA NATURE ET DE LA VIE ET SCIENCES DE LA TERRE ET DE L'UNIVERS
- Mougou, I. (2010). *Contribution à l'étude de la sensibilité de l'olivier à la maladie de la tuberculose (Pseudomonas stavenoi)*.
- Ministère du Développement et de la Coopération Internationale
- Mishra A-K. et Dubey N-K. (1994). . Evaluation of Some Essential Oils for Their Toxicity against Fungi Causing Deterioration of Stored Food Commodities. . *Applied and Microbiology.* , 60 (4): 1101-1105
- Moussaoui.R, Siziani.D, Youyou.A, & ., S. P. ((2010).). Antioxydant effect of phenolic compounds recovered from olive mill wastewater of chemlal variety cultivated in Kabylia (Algeria) on the oxidative stability of virgin olive oil. *Journal of food, Agriculture & Environnement.* , Vol 8 (3& 4): 86-89.
- Mueller, J. a. (1941). . *Proc. Soc. Exp. Biol. Med.*, 48: 330-333. .
- Mujica M-V., G. M. (2009). Importance of the extraction method in the quantification of total phenolic compounds in *Phaseolus vulgaris* L.Venzuela. . *Interciencia* , Vol. 34, N° 9: 650-654.
- N, G. K. (2007). *Etude de la diversité génétique de l'olivier cultivé en Tunisie : Approche pomologique, chimique et moléculaire. Thèse de doctorat en Sciences*.
- Nelson, P., & Toussoun, T. a. (1983). *Fusarium species: An Illustrated Manual for*

Identification. Pennsylvania State University Press., University Park. , . 193pp.

▪ Nmila,R,R.,Gross,M.Rchid,M.Manteghetti,P.P etit,M.Tijane and G.Ribes,2000.Insulin tropic effect of citrullus colocynthis fruits extracts.Planta Med.,66:418-423

▪ Office National de l'Huile, O. (2010). 10, Av. Mohamed V.

▪ Ownagh A., H. A. (2010). , 2010. Antifungal effects of thyme, agastache and satureja essential oils on *Aspergillus fumigatus*, *Aspergillus flavus* and *Fusarium solani*. . *Veterinary Research Forum* , 2: 99-105.

▪ PRABAVATHY, V.R., MATHIVANAN, N., SAGADEVAN, E., MURUGESAN, K.& LALITHAKUMAN, D., 2006. Self-fusion of protoplasts enhances chitinase production and biocontrol activity in *Trichoderma harzianum*. *Bioresour. Technol* 97, 2330-2334.

▪ P, L. (2003). Phytopathologie: bases moléculaires de biologie des pathosystèmes et fondement des stratégies de lutte. De Boeck & Presses Agronomiques de Gembloux (Eds.), Brussels, Belgium, p. 149-167.

▪ Pinto and Morais, 2000; Trapero and Blanco, 1999; USEPA, 1996 .Options Méditerranéennes, A, no. 106, 2013

▪ P.M., H. S. (2000). The role of polyphenols in terrestrial ecosystem nutrient cycling. *Trends in Ecology and Evolution*. , 15: 238-243.

▪ Parekh J. et Chanda S. V. 2007. *In vitro* antimicrobial activity and phytochemical analysis of some Indian medicinal plant. *Turkish journal of biology*., 31 : 53-58

▪ PETERSON, J.D.M., 1998. Flavonoids: dietary occurrence and biochemical activity. *Nutrition Res* 18, 1995-2018.

▪ Qazan WSH, Almasad MM and Daradka H. Short and long effects of *Citrullus colocynthis* L. on reproductive system and fertility in female Spague-Dawley rats. *Pak J Biol Sci*.2007;10(16):2699-2703

▪ Ramanatsoa, M. L. (2009). *efficacité des fongicides de remplacement dans la lutte contre les agents de la fonte semis attaquant les boutures de caféier*.

▪ Rahuman AA and Venkatesan P. Larvicidal efficacy of five cucurbitaceous plant leaf extracts against mosquito species. *Parasitol Res*. 2008;103(1):133-139.

▪ Ranalli.A. (1991). The effluent from olive mills: Proposals for re-use and purification with reference to Italian legislation. . *Olivae* , 37, 30-39.

▪ Rhouma, A. (2012). Principales maladies de l'olivier en Tunisie.

▪ Rota M. C., Herrera A., Martinez R. M., Sotomayor J. A. et Jordán M. J. 2008. Antimicrobial activity and chemical composition of *Thymus vulgaris*, *Thymus zygis* and *Thymus hyemalis* essential oils. *Food control*., 19 : 681-687.

▪ Ranalli.A. (1991). The effluent from olive mills: Proposals for re-use and purification with reference to Italian legislation. . *Olivae* , 37, 30-39.

▪ Roy,R.K.,M.Thakur and V.K.Dixit,2008.Effect of citrullus colocynthus on hair growth in albino rats .*Pharmaceutical Biol.*,45:739-744

▪ Ramanathan,T.,S.Gurudeeban and K.Satyani,2010.antioxidant and radical scavenging activity of citrullus .*Int Rapid Nutraceuticals*,1:37-37

▪ Rahbar AR and Nabipour I. The hypolipidemic effect of *Citrullus colocynthis* on patients with hyperlipidemia. *Pak J Biol Sci*. 2010; 13(24):1202-1207

▪ Rhouma A., H. B. (2008). *Antimicrobienne des activités de Pistacia et Schinus species against some plant pathogenic fungi and bacteria. Scientific research and essay*.

▪ Rhouma, A. (2012). Principales maladies de l'olivier en Tunisie.

▪ Saba AB and Oridupa AO. Search for a novel antioxidant, anti-inflammatory/ analgesic or anti-proliferative drug: Cucurbitacins hold the ace. *Journal of Medicinal Plants Research*. 2010;4(25): 2821-2826.

▪ S. Krid1, , A. (2010). Pseudomonas savastanoi endophytic bacteria in olive tree knots and antagonistic potential of strains of pseudomonas fluorescens and bacillus subtilis.

▪ Subrahmanyam M., H. A.-G. (2001). Antibacterial Activity Of Honey On Bacteria Isolated From Wounds . *Annals of Burns and Fire Disasters* , XIV (I).

▪ S. Shibko, P. Koivistoinen, C.A. Tratnyek, A.R. Newhall, L. Friedman.,1976. A method for sequential quantitative separation and determination of protein, RNA, DNA, lipid, and glycogen from a single rat liver homogenate or from a subcellular fraction *Analytical Biochemistry*, Volume 19, Issue 3, June 1967, Pages 514-528

▪ Satish S., R. M.-P.-C.-A. (2010). In vitro evaluation of the antifungal potentiality of *Polyalthia longifolia* against some sorghum grain moulds. . *Journal of Agricultural Technology*. , Vol.6(1): 135-150.

▪ S. Krid1, A. R. (2010). Pseudomonas savastanoi endophytic bacteria in olive tree knots and antagonistic potential of strains of pseudomonas fluorescens and bacillus subtilis. *Journal of plant pathology* , 92 (2), 335-341.

▪ SEGER, C.H., STRUM, S., MAIR, M.E., ELLMERER, E.P. & STUPPNER, H.,2005. ¹H and ¹³C NMR signal assignment of cucurbitacin

derivatives from *Citrullus colocynthis* (L) Scharder and *Ecballium elaterium* L. (Cucurbitaceae). *Magn. Reson. Chem* 43, 489-491.

▪ Sarni-Manchado P, C. V. (2006). Les polyphénols en agroalimentaire, Ed. Lavoisier (Tec & Doc), Paris, 300-398.

▪ Satish S., R. M.-P.-C.-A. (2010). In vitro evaluation of the antifungal potentiality of *Polyalthia longifolia* against some sorghum grain moulds. *Journal of Agricultural Technology*, Vol.6(1): 135-150.

▪ SMAOUI, S. (2010). *Purification et Caractérisation de Biomolécules à partir de microorganismes*.

▪ Solis-Pereira S., E. F.-T.-G.-G. (1993). Effect of different carbon sources on the synthesis of pectinase by *Aspergillus niger* in submerged and solid state fermentation. *Appl. Microbiol. Biotechnol.*, 39: 36-41.

▪ Spooner-Hart, V. S. (2009). Anthracnose and Cercosporiose on olives in Australia: an update. *Australian & New Zealand Olivegrower & Processor*, 31.

▪ Subrahmanyam M., H. A.-G. (2001). Antibacterial Activity Of Honey On Bacteria Isolated From Wounds. *Annals of Burns and Fire Disasters*, XIV (I).

▪ SAWAYA, W.N., DAGHIR, N.J. & KHALIL, J.K., 1986. *Citrullus colocynthis* seeds as a potential source of protein for food and feed. *Journal-of Agricultural and Food Chemistry* 34(2), 285-88.

▪ Solis-Pereira S., E. F.-T.-G.-G. (1993). Effect of different carbon sources on the synthesis of pectinase by *Aspergillus niger* in submerged and solid state fermentation. *Appl. Microbiol. Biotechnol.*, 39: 36-41.

▪ Simmons, J.G., E.N.T. AkobUNDU and J.P. Cherry, 1982. Chemical function and Nutritional properties of Egusi seed protein products. *J. Food Sci.* 47/829-835

▪ .

▪ Tiwari B-K., V. V.-P.-P. (2009). Application of natural antimicrobials for food preservation. *Journal of Agricultural and Food Chemistry*, 57: 5987-6000.

▪ Thakore, Y. (2006). The biopesticide market for global agricultural use. *Industrial Biotechnology*. 2: 194-208.

▪ thèse de DUTERTRE Julie Marie-Josèphe 2011 : DIPLOME d'ETAT de DOCTEUR EN MEDECINE : Enquête prospective au sein de la population consultant dans les cabinets de médecine générale sur l'île de la Réunion : à propos des plantes médicinales, utilisation, effets, innocuité et lien avec le médecin généraliste. Université Bordeaux 2 - Victor Segalen U.F.R DES SCIENCES MEDICALES

▪ Tiwari B-K., V. V.-P.-P. (2009). Application of natural antimicrobials for food preservation. *Journal of Agricultural and Food Chemistry*, 57: 5987-6000.

▪ Triki M. A., A. H. (2006). *Premières observations de Verticillium dahliae sur Olivier en Tunisie*. Bulletin OEPP, 36 (1): 69 -71.

▪ Triki, A. R. (2008). *First report of Pseudocercospora cladosporioides, the causal agent of Cercospora leaf spot of olive trees, in Tunisia*. *Phytopathol. Mediterr.* 47, 262-265.

▪ Ünal, M. ((1994).). Polyphénol, o-diphénols et acides phénoliques totaux dans les grignons d'olive et les margines. *Olivae*, 51:34-35.

▪ Vazequez R.A., M. D. (1974). *Compentes fenolicos de la aceituna. II*.

▪ Werner, H. (Mars 2009). *Rhizoctonia, cause de pourritures racinaires sur de nombreuses cultures*.

▪ Widden, P. (s.d.). Competition and the fungal community. Dans: The mycota IV. Environmental and Microbiological (Esser, K, Wicklow, D.T., Svdstrvm, B.E.). Springer-Verlag, Berlin, Allemagne. 1994, Pp. 135-147.

▪ Yen T-B. et Chang S-T., 2008. Synergistic effects of cinnamaldehyde in combination with eugenol against wood decay fungi. *Bioresource Technology* 99: 232-236.

▪ Yrjönen T., 2004. Extraction and Planar Chromatographic Separation Techniques in the Analysis of Natural Products. *Conference Room 513 at Viikki Infocentre (Viikinkaari 11), Faculty of Pharmacy of the University of Helsinki*: 64.

▪ Yuri. (2012). *Adhesive Tape Preparation for Examination of Fungi*.

▪ Z., M. (2006). Etude du pouvoir antimicrobien et antioxydant des huiles essentielles et flavonoïdes de quelques plantes de la région de Tlemcen. Thèse magistère. Université de Tlemcen:

▪ Zarrin M., A. N.-S. (2010). In vitro antifungal activity of *satureja khuzestanica* jamzad against *Cryptococcus neoformans*. *Pak. J. Med. Sci*, 26 (4): 880-882.

▪ ZIYYAT, A., LEGSSYER, A., MEKHFI, H., DASSOULI., SERHROUCHNI, M. & BENJELLOUN, W., 1997. Phytotherapy of hypertension and diabetes in oriental Morocco. *Journal of Ethnopharmacology* 58, 45-54.

• ZORO, A.B.I., KOFFI, K.K. & DJÈ, Y., 2003. Caractérisation botanique et agronomique de trois espèces de cucurbitées consommées en sauce en Afrique de l'Ouest : *Citrullus* sp., *Cucumeropsis mannii* Naudin et *Lagenariasiceraria* (Molina) Standl. *Biotechnol. Agron. Soc. Environ* 7 (4), 189-199