

# The GEMS (Gravity-EM Super) Unification Theory : the Unification of the Four Forces of Nature, Prediction of New 21 MeV and 22 MeV Particles, and Correspondence with Electro-Weak Theory

John E. Brandenburg PhD , Morningstar Applied Physics LLC  
Madison USA

**Abstract**—This manuscript presents some recent results of the GEMS (Gravity-Electro-Magnetism Super) Unification theory which unifies the four forces of nature. The two long range forces Gravity and Electro-Magnetism are first unified, and out of this unification also proceeds the unification of the short range Weak and Strong Nuclear Forces. They are unified under the two postulates that: 1. Gravity fields are an array of electromagnetic cells and 2. The separate appearance of Gravity and EM fields from each other is correlated with the separation of protons and electrons from each other as they emerge from the Planck scale with the appearance of a compact or hidden dimension. In the Standard Model all massive particles are charged and move freely at short distances and even photons spend time as charged particles. The quark-electron split occurs based on the asymmetry in dimensionality between space and time, with 3 quark colors representing space and the electron representing time. The proton, with its three interior quarks is born with the same effective radius as the electron- the electrostatic classical radius, with the gluon-photon separation also occurring. The theory produces the value of G: the Newton gravitation constant, and the proton mass accurately from the Planck scale with no free parameters. The theory produces the values of the masses, charges and spins for the pions of the Strong Force and the W and Z bosons of the Weak Force as quantum Mie scatterings off the compact dimension structures associated with the proton and electron masses. The Higgs Boson mass follows from similar formalism. The GEM theory extends the Standard Model to include Gravitation and a detailed correspondence is shown with Electro-Weak theory. The theory predicts a short lived, neutral spin 0 particle will be found at rest mass-energy approximately 22 MeV, and another at 21 MeV and that a basic GEMS parameter  $\sigma = 42.8503$ , occurs throughout the Standard Model in ratios of particle masses.

**Keywords**—GEM Super Unification Gravitation Electromagnetism quarks, Radion, Strong Force, Weak Force, Pions, W and Z Bosons, Higgs Boson ,gluons, New Particles 22MeV

## I. INTRODUCTION

The GEMS (Gravity-Electro-Magnetism Super) Unification theory, unifies all Four Forces of nature: the short-range Strong and Weak Nuclear Forces and the long-range Gravity and electromagnetic Forces for the first time in history. [1-6] Basically, it is a classical theory extended to subatomic limits, with appropriate quantum conditions on those limits. GEMS is basically a combination of the Kaluza-Klein hidden dimension approach to EM gravity unification and the Sakharov approach wherein space-time is electromagnetic in character. This leads to a definition of the spacetime metric as a normalized portion of the EM stress tensor.

(1)

$$g_{\alpha\beta} = \frac{4F_{\alpha}^{\gamma} F_{\gamma\beta}}{F_{\mu\gamma} F^{\mu\gamma}}$$

where  $F_{\gamma\sigma}$  is the Faraday tensor of relativistic EM.

The theory first unifies Gravity and Electromagnetism, the two classical long range forces of nature, with the constraint that the universe produce a hydrogen-rich Big-Bang. That is, a universe is postulated that from its beginnings is dominated by two charged particles protons and electrons, forming the source terms for the two long range forces, gravity and electromagnetism. The two forces, which began as a unified “super-force” correlated with a particle-antiparticle pair which evolves into the proton-electron pair as the universe expands from the Planck Scale. The creation of this correlated particle-force field pair, a strong and weak long range force, and heavy and light charged particle is controlled by the “deployment” of a hidden 5<sup>th</sup> dimension from initial size of a Planck Length  $r_P = (G\hbar/c^3)^{1/2}$ , to fully deployed size  $r_o$  of approximately 1/43 a classical electron radius:

$$r_o = \frac{e^2}{m_o c^2} \quad (2)$$

$$E\ell / c = N\sigma h \quad (5)$$

$$\ell = (r_o + \alpha r_e) \quad (6)$$

where  $m_o c^2 = (m_p m_e)^{1/2} c^2 = 21.897 \text{ MeV}$ , where  $m_p$  and  $m_e$  are the masses of the proton and electron respectively. This leads to the prediction of the discovery of a new particle, called the Morningstar or  $M^*$  particle, [4-6] representing the Kaluza-Klein 5<sup>th</sup> dimension, at mass-energy 22.897 MeV, with charge and spin of zero. This particle is in the seemingly region of mass energy states between the Muon and the electron.

$$m_o c^2 = \frac{e^2}{r_o} = 21.897 \text{ MeV} \quad (3)$$

This particle represents the 5<sup>th</sup> dimension that unifies gravity and EM.

It is seen that  $r_o = \sigma r_e$ . where  $\sigma$  is the central parameter of the GEMS theory :

$$\sigma = \sqrt{\frac{m_p}{m_e}} \quad (4)$$

It is found that when the 5<sup>th</sup> dimension and classical radii of particles form, these spherical structures behave as “branes” interacting with the surrounding quantum ZPF (Zero-Point Fluctuation) and to support quantum oscillations and resonances.[4-6] The lowest energy resonance is that associated with the electrostatic radius of the electron, a radius which is shared with the proton, ( the neutron scattering radius) and leads to the Strong Force carried by Pions (  $\pi$  mesons). The higher energy resonance is associated with the classical electro-static radius of the proton, and creates the Weak Force, mediated by the W and Z Bosons. Thus, in the GEMS unification theory, the splitting of the proton-electron pair into two distinct masses, is mirrored the creation of two separate long-range forces of distinct strengths, gravity and electromagnetism, and also the splitting of the two subatomic ranged forces into two forces of distinct strength, the Strong and Weak. The Higgs boson appears in this theory as an excitation of the surface of the 5<sup>th</sup> dimension, but its most fundamental manifestation is as a particle called  $M^*$  at 21.8965 MeV. Other particles that are quantum resonances or Quantum Mie Scatterings off the proton, electron or hidden dimension itself are found by the following formulas similar to the following [6]:

A family of particles, representing perturbed resonances of this 5<sup>th</sup> dimension should exist. Like any resonant oscillator, it can be perturbed by the presence of a nearby oscillator of different frequency.

One perturbed state has already been found at 16.7 MeV, [7] and is interpreted in the GEMS theory as a EM perturbed state caused by the near-lying electron state:

$$E = M^* c^2 / (1 + \alpha\sigma) = 16.6806 \text{ MeV} \quad (7)$$

Which is approximately what has been found. This indicates a family of mass states may be found in the region of 22-17 MeV. Following previous patterns, another state should be found at

$$E = M^* c^2 / (1 + 5\alpha) = 21.126 \text{ MeV} \quad (8)$$

Which we will call  $M^{**}$

In the rest of this article a brief synopsis of the GEMS theory will be given, with a summary of its results and predictions. A detailed correspondence between GEMS and Electro-Weak theory has also been found and will also be presented.

## II. THE POSTULATES, MODELS, AND BASIC RESULTS OF THE GEM THEORY

The following explains how the basic concepts of the GEMS theory are turned into models and their basic results. The GEMS theory is a combination of the Kaluza-Klein [8] and Sahkarov [9] approaches to EM-Gravity unification, where a hidden 5<sup>th</sup> dimension allows separate EM and gravity fields, but that spacetime itself is electromagnetic and consists of the quantum ZPF. Under the conjecture of Dr. Alfred Luhen, (Private Communication) ‘one cannot create mass without creating gravity.’ The Higgs Boson thus cannot exist and generate mass outside the context of General Relativity. In the GEMs context the Higgs scalar field occurs in Kaluza-Klien theory as the Radion scalar field so that both gravity, EM-mass energy as gravity source term, and particle charges and masses are born together with the Radion field.

### A. Gravity Fields and Spacetime as Electrodynamics

The first basic postulate of the GEM theory is that gravity fields can be synthesized as arrays of  $E \times B$

drifts familiar from plasma physics. The concept for a synthesis of a gravity field from electromagnetism was the outgrowth of the effort to achieve controlled thermonuclear fusion, most specifically the magnetic confinement of plasmas for fusion. As part of this effort the motion of charged particles in magnetic and electric fields was carefully studied and an effect called an "E-cross-B drift" or ExB drift [10], was identified.




Fig. 1. The ExB drift caused by crossed electric and magnetic fields affects all charged particles identically and in non-uniform E fields, but uniform B fields, can cause acceleration.

This effect is remarkable in that it affects all charged particles identically regardless of charge or mass. We can derive this model of a gravity field simply by first assuming uniform E and B fields at right angles to each other, as in Figure 1, for example,  $E_x$  and  $B_z$  in the x and z directions respectively. We have then for motion of a charged particle in the x and y directions or  $r, \theta$ , using esu units:

$$m \frac{\partial V_x}{\partial t} = qE_x + \frac{V_y}{c} B_z \quad (9)$$

$$m \frac{\partial V_y}{\partial t} = qE_y - \frac{V_x}{c} B_z \quad (10)$$

Where we have included an  $E_y$  for a curvilinear E field. We can solve this by assuming a velocity function of two parts, in x and y coordinates. Here we make the simplification that  $E_x \gg E_y$ , i.e. a particle at the center of the region between the two plates in Figure 1.

$$V = V_{osc} + V_d \quad (11)$$

$$V_d = \frac{cE_x}{B_z} \quad (12)$$

in the y direction with the definitions

$$V_{osc,y} = V_{\perp} (\sin \omega_c t) \quad (13)$$

$$V_{osc,x} = V_{\perp} (\cos \omega_c t) \quad (14)$$

Where  $V_{\perp}$  is assumed to be a constant with  $V_{\perp} \leq V_d$  and we have defined

$$\omega_c = eB/mc \quad (15)$$

Note this drift velocity shown in Eq. 12 is independent of charge and mass.

If we leave the magnetic field uniform and vary the E field at right angles to its direction, in the direction of the drift, the particle will experience an acceleration in the direction of its ExB drift in the y direction:

$$\frac{\partial V_d}{\partial t} = \frac{c^2 E_x}{B_z^2} \frac{\partial E_x}{\partial y} = \frac{\partial \phi}{\partial y} \quad (16)$$

$$\phi = \frac{1}{2} \frac{E_x^2}{B_z^2} c^2 \quad (17)$$

This is easily confirmed by a particle simulation where an electron and a 'heavy positron' of positive charge but 10x the mass of electron are released in uniform magnetic field but between two plates set at an angle between each other, as seen in Figure 2.




Fig. 2. A simulation of an EM-synthetic gravity field with the trajectories of an electron and a "heavy positron" of 10x an electron mass are seen.

The gyro-motion radius  $a_0$  of the particles seen here vanishes in the limit of very strong magnetic fields ( $B_z \rightarrow \infty$ ) thought to be present in the vacuum due to the quantum ZPF (Zero Point Fluctuation)

whereas the gravity produced velocity is  $cE_x/B_z$  is much less than light for ordinary gravity fields.

$$a_o = \frac{E_x}{B_z} \frac{mc^2}{e} \rightarrow 0 \quad (18)$$

We have found this physical model of gravity fields as being composed of locally uniform magnetic and varying electric fields. Flat spacetime then, we can conceive of as being composed of uniform magnetic and electric fields. But the vacuum is observed to be massless, or even to have a tiny negative mass density. How then, is the vacuum full of powerful fields to create an ExB drift array to create gravity, but yet has no mass density? To be consistent with GR, the mass density of the very E and B fields causing gravity must be considered as a source of gravity. This problem is not unique to the GEM theory but is a pressing problem for any theory of a quantum vacuum.

Einstein discovered the ZPF (Zero Point Fluctuation) in 1910, showing that as a consequence of the Heisenberg Uncertainty principle the vacuum itself must be populated with EM modes. The physical presence of these modes can be seen in the existence of the Casimir Effect. However, the fact that these modes do not create an observable mass-energy density in the vacuum is one of the great mysteries of physics. This problem was considered by the great Russian physicist Yakov Zeldovich [11] who argued that a ZPF mass density term would appear as a Cosmological Constant term, allowed by General Relativity, and that another such term existed to cancel the ZPF term. The Zeldovich Cancellation term would then be required for a massless vacuum that we experience. Here we have the basic field equation for GR including the Cosmological Constant  $\Lambda$ :

$$R_{\nu\mu} - \frac{1}{2} g_{\nu\mu} R = \frac{8\pi G}{c^4} T_{\nu\mu} - \Lambda g_{\nu\mu} \quad (19a)$$

$$\frac{8\pi G}{c^4} T_{\nu\mu} - \Lambda g_{\nu\mu} = 0 \quad (19b)$$

GEM theory is an alloy of the concepts of Sahkarov [9], in gravity's relationship to the EM ZPF, and the Kaluza-Klein theory [8] of EM-gravity unification, and its relationship to a hidden 5<sup>th</sup> dimension. To see this we begin with the Hilbert action principle in 4 spacetime dimensions with a zero cosmological constant.

$$W = (16\pi G)^{-1} \int R \sqrt{-g} dx^4 \quad (20)$$

where R is the Curvature Scalar. Finding the extremum of this action leads to the vacuum gravity equations with canceled ZPF EM fields.

$$R_{\nu\mu} - \frac{1}{2} g_{\nu\mu} R = 0 \quad (21)$$

Sahkarov interpreted the integrand as a real energy density. He equated this energy density to a perturbed quantum EM ground state spectrum of ZPF (Zero Point Fluctuation) due to the Heisenberg Uncertainty principle applied to the vacuum EM field. The zeroth-order ZPF is assumed to vanish due to a canceling cosmological constant term proposed by Zeldovich. The "Zeldovich Cancellation" ensures that only the perturbations due to curved space cause the effect of the ZPF to appear. Sahkarov calculated the perturbed part of the ZPF due to spacetime curvature. He then derived a formula for G in terms of an integral over the perturbed ZPF:

$$W = G^{-1} \cong \frac{\hbar}{2c^5} \int_0^{\omega_P^*} \omega d\omega = \frac{\hbar \omega_P^2}{c^5} \quad (22)$$

$$G \cong \frac{c^3 r_P^2}{\hbar} = \frac{c^4}{r_P^2 T_o} \quad (23)$$

where  $\omega_P$  is the Planck frequency  $c/r_P$ , where  $r_P = (G\hbar/c^3)^{1/2}$  and the energy density  $T_o = \hbar c/r_P^4$  is the Planck scale energy density. This is consistent with a physical model of gravity forces as due to imbalances of the EM Poynting vector,  $S = cExB/4\pi$  ( in esu) or a radiation pressure  $P = \langle S \rangle / c$ . The second example of radiation pressure or Poynting vector acting on particles in a box whose walls absorb and emit radiation is shown in Figure 3. In Figure 3, the left figure shows hot-bright particles in a dark-cold enclosure, the right figure shows cold-dark particles in a hot-bright enclosure. Mutual radiation pressure forces are shown by block arrows.




Fig.3. Radiation Pressure Affecting Particles in an Enclosure. Left: Two hot ideal radiators in a cold box repel each other by mutual radiation pressure. Right : Two cold ideal radiators in a hot box attract each other due to mutual shadowing.

As was shown in the first section an ExB or Poynting drift field, with constant B and E growing stronger in the direction of the drift, can produce

gravitational-like acceleration of charged particles of all charges and masses, as shown in Figure 1. The Sakharov model for the gravitational force is basically that of a radiation pressure Poynting field produced by non-uniformities in the ZPF and is successful in the sense that is self-consistent (see Figure 3). It is understandable that Sakharov would arrive at this physical model for gravity, since he worked on the Soviet Hydrogen Bomb where radiation pressure is crucial. We can derive the same idea, in relativistic-covariant form, from the expressions in the first GEM article [2], where the zeroth-order ZPF stress energy was caused to vanish. That is we will explain the Zeldovich Cancellation as EM-gravity unification physics.

The following equations show this theory in covariant form. It can be seen that if the metric tensor for gravity is written as a normalized first part of the EM momentum-stress tensor:

However, if the fundamental structure of spacetime is electro-magnetic we can write the metric tensor as an electromagnetic tensor[2] :

$$g_{\alpha\beta} = \frac{4F_{\alpha}^{\gamma} F_{\gamma\beta}}{F_{\mu\nu} F^{\mu\nu}} \quad (24)$$

For the case of statistically uniform isotropic vacuum fields it is easy to see that the elements of the gradient of the metric will vanish.

When this expression is used, the EM stress tensor for the ZPF can be made to vanish as shown in the first article on the GEM theory [2].

$$T_{\alpha\beta} = F_{\alpha}^{\gamma} F_{\gamma\beta} - g_{\alpha\beta} \frac{F_{\mu\nu} F^{\mu\nu}}{4} = 0$$

Here we assume a model of spacetime containing adjacent regions of strong E or B fields. The particles however, travel as wave packets and sample a volume swept out by a wave-front, thus they see an average spacetime. An average over volume so that  $\langle B^2 \rangle = \langle E^2 \rangle$  and  $\langle E \cdot B \rangle = 0$  results in a volume average of two metric forms one dominated by electric flux, for instance, in its local direction  $E_y$

$$g_{\alpha\beta} = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (26)$$

And another, in an adjacent region, by magnetic flux also in  $B_y$

$$g_{\alpha\beta} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -2 \end{pmatrix} \quad (27)$$

Upon volume average, assuming large scale isotropy, we recover the familiar Lorentzian flat space metric.

$$\langle g_{\alpha\beta} \rangle = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad (28)$$

Using the observation that, for nearly flat spacetime, gravity fields and their potentials are linearly additive, we can derive the effective gravity potential for the ExB drift model of gravity assuming the EM form of the metric tensor required for self-censorship. We then find for the upper left diagonal element of the metric tensor:  $g_{00}$ , and from it the effective Newtonian gravity potential.

(29)

$$g_{\alpha\beta} = 2[B^2 - E^2]^{-1} \begin{pmatrix} -E^2 & 0 & 0 & 0 \\ 0 & E_x^2 - B_y^2 - B_z^2 & 0 & 0 \\ 0 & 0 & E_y^2 - B_x^2 - B_z^2 & 0 \\ 0 & 0 & 0 & E_z^2 - B_x^2 - B_y^2 \end{pmatrix}$$

We have then for perturbing fields and a gravity potential in terms of an  $E \times B$  drift model of gravity that is valid for both DC and oscillating E fields, where charged particles are accelerated into the strongest

part of the perturbing E field. How then does the Newtonian gravity potential between charged particles come about? We begin with the expression for a gravity potential in terms of E and B fields in the vacuum, where  $V_D$  is the particle drift velocity in the crossed E and B fields:

$$\langle g_{00} \rangle = 1 + 2\phi/c^2 \cong \frac{-E^2}{B^2 - E^2}, \quad E^2 = E_x^2 \quad (30)$$

We obtain from GEM Metric tensor to first order in  $E_x/B_z \ll 1$  and averaging with a flat metric.

$$-1 - 2\phi/c^2 = -\frac{E_x^2}{B^2} \quad (31)$$

As was first proposed by Puthoff [12], it can be shown [13] that point charges floating in a ZPF will create an interference pattern  $E^2$  between their scattered  $1/r$  radiation E fields and the impinging ZPF E fields, leading to a Newtonian potential around each particle. It was also pointed out by Puthoff that under the Standard Model all fundamental massive particles are charged and move freely at short distances, consistent with electrodynamic gravity. QED ensures that even photons spend part of their time as charged particles and are thus subject to electrodynamic gravity [5]. Using the metric formulation of Eq. 24 and spatial averaging a full Schwarzschild Metric:  $g_{rr} = 1/(1 - 2GM/c^2 r)$ , arises statistically around each charged particle [13, 5].

### B. Particles From the Vacuum: The GEM Concept

We have the vacuum quantities associated with the Planck scale, the Planck length, the Planck mass, and the Planck charge, respectively:

$$r_p = \sqrt{\frac{G\hbar}{c^3}} \quad (32a)$$

$$M_p = \sqrt{\frac{\hbar c}{G}} \quad (32b)$$

$$q_p = \sqrt{\hbar c} \quad (32c)$$

The simplest result then would use the vacuum derived Planck charge  $q_p$  as the length of the path in the 5<sup>th</sup> dimension. Using this we could obtain the proton mass as the simplest result.

We must now consider other constraints to such a theory. Nothing, especially the cosmos itself, is by definition simple. In particular, the appearance of one particle does not increase entropy in the universe, and entropy requires complexity. Also, we must consider that a charged particle cannot simply pop out of the vacuum without violating the electromagnetic constraint of charge neutrality. So the same simple process of a path integral allowing the appearance of a proton must also allow the appearance of an electron to balance it and to maximize entropy.

Therefore, we must have the proton appear as part of a system that includes the electron, so that hydrogen results:

$$q_p = -q_e \quad (33a)$$

$$q_p = e, \quad -e = q_e \quad (33b)$$

Another constraint occurs because the path length in the vacuum that cannot be simply a distance, but must be a spacetime interval. In the vacuum state all particles must be masses and move at the speed of light and have a spacetime interval of zero:

$$r_o^2 = (x_o^2 + y_o^2 + z_o^2) \quad (34a)$$

$$r_o^2 - c^2 t_o^2 = 0 \quad (34b)$$

It is seen that the appearance of the new hidden dimension occurs in a form analogous to the splitting of a canceling charge pair of particles from the vacuum, by splitting of a quantized light-like, or vacuum, space-time interval of length zero. In the GEM theory the hidden dimension size, where the hidden dimension can mix with the non-hidden dimensions, is the quantized particle size. The hidden dimension quantities are thus able to mix with the normal spacetime quantities because they are similar at smaller scales. This will lead to, as we experience them, two particle types. One is associated with the time-like portion of the constrained interval, leading to a one-dimensional scalar character, an electron, and another of equal size with a space-like vector character having three constrained sub-dimensions, a proton. The gravitation constant  $G$ , functions in the vacuum as the "interpreter" of charge into either mass or distance. Thus, ironically, charge and mass, the source terms for EM and gravity, are unified already in the vacuum quantity  $G$ , which has units of charge to mass ratio squared in the esu system used here.

$$q\sqrt{G/c^4} = r_o \quad (35a)$$

$$r_o^2 = (G/c^4)(q_x^2 + q_y^2 + q_z^2) \quad (35b)$$

$$r_o^2 = c^2 t_o^2 = (G/c^4)q_t^2 \quad (35c)$$

Therefore, the quantized vacuum scale length, the Planck length, gives birth to a quantized larger scale hidden dimension. Because the quantized hidden dimension is an image of macroscopic space-time in a light-like interval, and its structure is part of a split "lightlike" spacetime where charge  $q$  is analogous to macroscopic dimensions as a length, we have charge conservation and interval conservation. We obtain

from these conditions the following constraints on the charges of the particles:

$$q_o = -q_4 = q_1 + q_2 + q_3 \quad (36a)$$

$$q_o^2 = q_4^2 = q_1^2 + q_2^2 + q_3^2 \quad (36b)$$

where the subscripts, 1,2,3,4 denote x,y,z,t the corresponding time or space dimensions in the unconstrained Cosmos.

Thus, the space-like portion of the split interval, the proton, has three sub-dimensions that we interpret as quarks or sub-charges, while the electron acts like a single entity. The gluon octet -photon separation also occurs at this juncture ( see Figure 4.)




Fig. 4. Formation of the gluon octet from the six-fold splitting of the photon pair. The eight faces of the double-tetrahedron correspond to the gluon octet.

This concept then makes the electron-quark family a reflection of the dimensional asymmetry of spacetime, with a scalar time and a three dimensional spatial dimensions. However, we have here specified the Planck length as the shared radius of the quarks and electrons. But the physics of the world depends on the much larger scale of subatomic particles. Here the electrostatic radius of the electron enters as the final "deployed" length for the electron. By the requirement The default radius upon "full deployment" of the hidden dimension and its physics would be the electron electrostatic radius  $r_e = \frac{1}{2} e^2/m_e c^2 = 1.4 \times 10^{-13} \text{cm}$  where we assume this radius will be shared by the quarks so that we have also  $r_c = \frac{1}{2} (q_1^2 + q_2^2 + q_3^2) / m_e c^2$  and this is the nucleon radius  $r_n \cong 1.4 \times 10^{-13} \text{cm}$  [14] determined by Strong force scattering, and also the charged pion Compton wavelength, considered the range of the Strong Force.

Thus, the conservation of vacuum interval and charge neutrality requires that the electron and proton share the same radius, as is approximately observed. This requires two conditions on the three dimensional array of quark charges

This concept of the electron and proton being born together explains both quark confinement and the absence of proton decay as geometric requirements, seeing as the proton must preserve its dimensionality

in quark space. That is, it is three dimensional and like any three dimensional object it cannot become an object of lower dimensionality.

This can only be satisfied by a SO(3) symmetry group, similar to the SU(2) symmetry group of Electroweak theory and is satisfied by the values

$$q_x = -\frac{1}{3}, q_y = q_z = \frac{2}{3} \quad (37)$$

This corresponds to the standard quark model. Thus, the GEM theory is actually compatible with the standard model.

In the GEM theory, the splitting apart of the proton and electron is correlated to the splitting apart of the gravity and EM forces. In the Standard Model context, this means that baryon and lepton number: **B** and **L** respectively, are not conserved but their difference (**B-L**) is conserved and the non-conservation of **B** and **L** separately occurs at the Planck scale, where gravity and EM unify. The appearance of charge and mass at the subatomic scale occurs with the appearance and deployment of the 5<sup>th</sup> dimension, which is slightly smaller than the EM cross-section of the electron. This means that, instead of subatomic particles being considered points, they must be treated as objects of definite size similar to the 5<sup>th</sup> dimension radius. This means that in the presence of the vacuum ZPF the structural sizes of the particles support resonances, and these resonances in-turn take on a quantum existence of their own.

In quantum electrodynamics, it is found that the sizes of various quantum objects can be understood as being created through orders of EM interaction. The Bohr radius of the hydrogen atom, and the Compton radius of the electron, for instance, can be found as the electron classical radius  $r_e = e^2/m_e c^2$  for instance, can be found as the  $1/\alpha^2$ , and  $1/\alpha$  respectively times the electron classical radius. However, the electrostatic radius for the electron is  $\frac{1}{2}$  the electron classical radius. This factor of  $\frac{1}{2}$  can be understood as the difference between monopole or "scalar" EM interactions, which cannot propagate farther than  $r_e$  and dipole "vector" EM waves which can propagate.

#### D. The Value of G and the Proton Mass From the Planck scale

The second GEM postulate is that Gravity and EM forces separate in correlated way with the appearance of electrons and protons from the Planck Scale. We can examine this by a Gedanken experiment where we squeeze a single atom of hydrogen in sphere until it becomes the size of a Planck radius and forms a

Blackhole. The Blackhole then evaporates via Hawking radiation [15] into a shower of gamma rays and particles and antiparticles and thus destroys the baryon and lepton number of the original electron ad proton.

Let us consider a “Gedanken” experiment [3] where a single atom of hydrogen is confined in a sphere whose size is shrunk continuously until it reaches approximately the radius of a Planck length  $r_P = (\hbar G/c^3)^{1/2}$ , (See Figure 5.) at this point the electron and proton making up the hydrogen will have long since ionized and increased in mass due to Heisenberg Uncertainty. The proton and electron will then form a Black Hole which will then undergo Hawking Evaporation [15] (Figure 5.) into a shower of photons, particles and their anti-particles. It is noticed that this evaporation will destroy the baryon and lepton number of the proton and electron, leaving only the quantum numbers of the vacuum. This is in accordance with the observation that many of the quantities we observe in the present day cosmos are “running constants” and change under radically smaller spacetime curvature, to merge eventually with Planck Scale quantities. Therefore, what we consider to be physical constants may be tied to specific range of scale-size for the radius of curvature of spacetime, and these physical quantities will change dramatically when the radius of curvature approaches the Planck Scale.




Fig. 5. A Gedanken experiment where 1. A single atom of hydrogen, a proton and an electron, is shrunk within a sphere 2. And ionizes. Finally it reaches the Planck size 3. And becomes a Black Hole, whereupon, 4., it undergoes Hawking Evaporation and becomes a cloud of gamma rays, matter and antimatter so that original hydrogen is lost.

In the previous chapter we were able to formulate gravity fields as electro-magnetic but this appeared to require a cellular nature for spacetime, in order to allow gravity fields, and spacetime itself, to be composed of regions of powerful electric and magnetic fields. At first, such a physical picture seems perfectly consistent with the concept of the Planck Scale, where spacetime is a foam of scale size equal to the Planck Length:  $r_P = (G\hbar/c^3)^{1/2}$ . However, at the Planck scale only a limited group of physical constants are possible and these do not include many of those constants that describe the universe we experience. We can imagine that in the primordial first instants of the Big Bang the entire universe was in a compressed state at the Planck Scale but that it

expanded from this scale to “deploy” a new larger scale that carried with it the physics of the cosmos we know. Therefore, in this chapter we must further quantify the concept of cellular spacetime to define a range of scale size for a cellular structure in spacetime that is distinct from the Planck Scale and represents an expanded scale that emerges from that primordial scale.

Thus, based on our Gedanken experiment, we consider that any cellular scale size in the vacuum is “fully deployed” to its proper size in the present cosmos, and helps determine its physics, but this scale size is crushed out of existence at the Planck scale, where hydrogen disappears. Accordingly, our Gedanken experiment to squeeze a proton-electron pair into the vacuum, also squeezes the cellular scale-size into the Planckian vacuum.

Let us assume however, in our thought experiment, that the wave functions of the proton and electron, carrying with them all their identifying quantum numbers have merged as the Black Hole forms at the Planck Scale, that is, the radius of spacetime local spacetime curvature  $r \rightarrow r_P$  before this happens. To model this behavior we will use a simple U(1) symmetry model for the proton and electron masses, considering that since all information disappears we will formulate the model only in terms of charge  $q$ , mass  $m$ , and mass ratio  $m_p/m_e = \sigma$ . Accordingly we have a simple U(1) mass model:

$$m = m_o \cos(\phi) + im_o \sin(\phi) \quad (38)$$

The U(1) symmetry is complex valued with real and imaginary mixed together. Particles with imaginary rest masses are tachyons, particles that move faster than light. The simplest physical interpretation we can make for such imaginary particles is that they are particles that have fallen inside the event horizon of a Black Hole, accelerating beyond the speed of light in the process and being out of communication with the real particles of the universe. This is important at the Planck scale because there particles appear out of the vacuum, form black holes and disappear, so that spacetime is effectively a “foam.” Foamy spacetime features Black Holes that are so closely packed that it is impossible to determine whether a particle is inside or outside an event horizon. Thus particles at the Planck scale can be physically represented as complex, half real and half imaginary, with masses satisfying a U(1) symmetry. So the Planck scale is completely chaotic, mixing imaginary masses with real ones.

Let us imagine that at the Planck scale everything becomes simple, the EM and gravity forces unify to one force obeying U(1) symmetry and lepton and baryon number also disappear, in fact everything disappears but “vacuum” quantities:  $G$ ,  $c$ , and  $\hbar$ , the Newton gravitation constant, the speed of light, and the rationalized Planck’s constant respectively. These determine the Planck Length:  $r_P = (G\hbar/c^3)^{1/2}$ . Planck mass  $M_P = (\hbar c/G)^{1/2}$ , and Planck charge  $q_P = (\hbar c)^{1/2}$



However, let us assume also that since EM forces still exist and enforce quantization of charge and the charge neutrality of the vacuum, so that quarks remain grouped in groups of 3 having one positive electron charge to cancel the charges of electrons. Thus, a plasma consisting of quarks and electrons occurs at the Planck scale, but protons are still identifiable as groups of quarks because the vacuum must be charge neutral.

Therefore, at the Planck scale we can have Planck masses of real and imaginary masses consisting of a quark-electron plasma which can still be represented as relativistic mass-dilated electrons and protons because of the requirement of charge neutrality.

On the other end of the spectrum of sizes we assume a "fully deployed" cellular scale of subatomic size, which we propose to be of the size range

$$r_o = \frac{e^2}{m_o c^2} \quad (39)$$

where  $m_o c^2 = (m_p m_e)^{1/2} c^2 = 21.897 \text{ MeV}$  so that the size scale is neutral between protons and electrons, and a size parameter which is determined entirely by low energy physics quantities. We will call this energy and size scale the "mesoscale" because it lies between the Planck Scale and the Cosmic Scale.

This is all based on the GEM postulate that baryon and lepton number disappear at the Planck scale coincidentally with the separate identity of Gravity and EM fields. The vacuum is thus as simple as possible at the Planck scale, only particles and anti-particles of Planck mass and charge exist there and gravity and EM are basically merged.

In contrast the appearance of the cellular scale size as the universe expands from the Planck scale represents the appearance of a new degree of freedom. This is similar to when a molecular layer evaporates from a surface and becomes a 3-dimensional gas. We will consider then, accordingly, that the expansion from the Planck Scale allows the appearance of a 5<sup>th</sup> dimension, represented by the appearance of a new scale size :  $r_o$ , which is the appearance of particles: electrons and protons with their classical radii. That is, the appearance of the 5<sup>th</sup> dimension allows the appearance of the mesoscale. The expansion of the universe from the Planck Scale thus allows a new 5<sup>th</sup> dimension, a new degree of freedom, of much larger scale size than the Planck Scale, to appear, and with it new physics. But how shall we include this into our U(1) mass model?

The angle  $\phi$ , we will consider, in this model, corresponds to charge state and is thus quantized as a canceling pair  $\pm\phi_o$ , even in the Planck Scale. However let us model the appearance of the fifth dimension by allowing this angle to become an imaginary rotation angle to give two real particle masses corresponding to an "up" quantum state and

"down" quantum state from the U(1) symmetry. Let us therefore assume a model of a scale dependent vacuum where the existence of a 5<sup>th</sup> dimension breaks the vacuum scale invariance. We now have for the mass model:

$$m = m_o \exp(\pm\phi_o) \quad (40)$$

Where  $\phi$  is a parameter such that  $\phi = 0$  at  $r \sim r_P$ . That is, near the Planck scale, when the 5<sup>th</sup> dimension does not exist and thus protons and electrons are identical. At the other extreme  $\phi = \phi_o$  when the 5<sup>th</sup> dimension is "fully deployed" and separate particle masses are generated at  $\phi_o$  from Eq. 40 as  $r \rightarrow r_o$ . This suggests a formula  $\phi_o \sim \ln \ln (r/r_P)$ , so that  $\phi_o$  very strongly near the Planck scale but varies very little at everyday scale. This mass function is of the form expected for dependence on the hidden Kaluza-Klein 5<sup>th</sup> dimension [8].

$$\sqrt{\frac{m_p}{m_e}} = \exp(\phi_o) = \sigma \quad (41)$$

Where  $\sigma$  is a mass asymmetry parameter, being the square root of the mass ratio of the electron to the proton.

Thus, even though mass symmetry is broken in terms of the new 5 space we experience, it is actually preserved in terms of a geometry involving the imaginary angles in the original U(1) symmetry. That is, the new particle dimension looks symmetric in the space of imaginary angle.

We require that this simple mass model give the behavior as  $m_o \rightarrow M_P$ ,  $\sigma \rightarrow 1$  as  $r/r_P \rightarrow 1$

To obtain a smooth transition to the Planck scale as curvature collapses to the Planck length the angle  $\phi_o$  must be dependent on curvature near the Planck length but very insensitive to it at larger curvatures, where the new fifth dimension is fully deployed. Based on the lack of observation of proton decay, lepton and baryon numbers are obviously strongly conserved. The simplest model to obtain this mixture of scale sensitivity with curvature  $r$  is for the rotation angle to have the dependence on our 5<sup>th</sup> dimensional deployment parameter

$$\phi \cong \ln(\sigma) \quad (42a)$$

$$\sigma \cong \ln(r/r_P) \quad (42b)$$

So that lepton and baryon numbers disappear, with  $\sigma \rightarrow 1$  as  $r \rightarrow r_P$

Therefore, in the GEM model, the separate appearance of proton and electron pairs from the vacuum is, like the separate appearance of EM and gravity forces, linked to the appearance and full development of the fifth dimension. The physical description of this new 5<sup>th</sup> dimension is that it comes into being at scale size that corresponds to the size of a particle classical radius  $r_o$ .

However, it is apparent Eq. 47 cannot be correct near  $r = r_p$  where  $\sigma \rightarrow 1$ , thus we must modify the formula slightly so that both the right and left side go to zero smoothly at  $r = r_p$  and  $\sigma = 1$ , where we assume  $\sigma$  goes to one with the vanishing of a small parameter  $\varepsilon \rightarrow 0$

$$\sigma = 1 + \varepsilon \dots \quad (43)$$

We rewrite Eq. 29:

$$\ln \left| \frac{r}{r_p} \right| = \sigma - \frac{1}{\sigma^2} \quad (44)$$

We now see that both sides of this expression go to zero as both quantities  $r/r_p$  and  $\sigma \rightarrow 1$  as they should. We have added the correction factor as second order in  $\sigma$ , that is  $\sigma^{-2} = m_e/m_p$  to be similar to the reduced mass correction of the conventional dynamics of the electron-proton system. Therefore, when the new 5<sup>th</sup> dimension is "fully deployed" we have for  $\sigma = 42.8503\dots$

$$\ln \left| \frac{r_o}{r_p} \right| = \sigma - \frac{1}{\sigma^2} \quad (45)$$

We note how both sides of this expression go to zero with leading order in  $\varepsilon$ , as  $r/r_p \rightarrow 1$ :

$$\ln \left| \frac{r}{r_p} \right| = \sigma - \frac{1}{\sigma^2} \cong 3\varepsilon \quad (46)$$

We must also correct the mass formula so that  $m_o = M_P$  at the Planck scale. So we must write, using the Planck charge  $q_P$ . We will assume that the normalized charge state assumes the role of determining mass  $q/e$  but that as we approach  $r = r_p$  that  $e \rightarrow q_P = (\hbar c)^{1/2}$  so that  $\alpha \rightarrow 1$  and also all masses approach the Planck mass  $m_o \rightarrow M_P$

$$m = m_o \exp\left(\frac{\pm q}{e} \ln \sigma\right) \quad (47)$$

This formula gives the observed mass difference between the electron and proton and also ensures that this difference disappears as  $r/r_p \rightarrow 1$ . However, not only mass the mass difference disappear but the mass  $m_o$  must undergo the process  $m_o \rightarrow M_P$ , as  $\sigma \rightarrow 1$  We therefore extending this formula, where normalized charge controls mass, to obtain

$$m_o \cong M_P \exp\left(\frac{q_P}{e} \ln \sigma\right) \quad (48)$$

Where this gives the proper limit as  $m_o \rightarrow M_P, \sigma \rightarrow 1$ .

However, we also require the condition, as  $m_o \rightarrow M_P$  that we must have the condition that  $r, r_p, m_o$  and  $M_P$  have the proper quantum relationship  $r_o = \hbar/(m_o c)$  so that near the Planck scale

$$\ln \left| \frac{m_o r_p}{M_P r_o} \right| = \ln \frac{m_o}{M_P} + \ln \frac{r_o}{r_p} = -3\varepsilon + 3\varepsilon = 0 \quad (49)$$

We obtain this behavior in  $\varepsilon$  for the  $m_o$  system by modifying Eq. 48, like we did the  $\sigma$  expression in Eq. 44, with a second order term  $\alpha$  to ensure the proper behavior for  $m_o$ , as  $\alpha^{-1/2}$  and  $\sigma \rightarrow 1$

$$m = M_P \exp(-(\alpha^{-1/2} + \alpha + 1) \ln \sigma) \exp\left(\frac{q}{e} \ln \sigma\right) \quad (50)$$

This requires, at normal spacetime curvature and charge state  $q/e = +1$  the expression for the proton mass, with  $M_P = 2.17645 \times 10^{-56} \text{g}$ :

$$m_p = M_P \exp(-(\alpha^{-1/2} + \alpha) \ln \sigma) = 1.6665 \times 10^{-24} \text{g} \quad (51)$$

This expression agrees with the observed rest mass of the proton  $1.67262 \times 10^{-24} \text{g}$ , to 3.6 parts per thousand and goes to the proper limit of  $m_p = M_P$  as  $\sigma \rightarrow 1$ .

We now return to primary expression relating normalized spacetime curvature to the mass ratio.

The expansion of the effective curvature to  $r_o$ , which we will term the "mesoscale" radius -since it is the range of scales of classical particle radii and lies between the Planck and Cosmic scales- then yields, by Eq. 50 the relation:

$$\ln \frac{r_o}{r_p} = \sigma - \frac{1}{\sigma^2} = 42.850\dots \quad (52)$$

If we examine the ratio of the Planck Length to the mesoscale radius, we discover it is also a quantum-EM normalized ratio of coupling constants between gravity and EM,

$$\frac{r_p}{r_o} = \sqrt{\frac{G m_o^2}{e^2 \alpha}} \quad (53)$$

This suggests that the gravitational interaction between two masses is mediated by the emission and absorption of EM photons. This is as we would expect if both EM and Gravity were both part of the same general phenomenon. The formula of Eq. 53 can be inverted to find an accurate expression for the gravitation constant.

We thus obtain for the gravity constant, using the measured value of the proton electron mass ratio, to first order:

$$G = \frac{e^2}{m_p m_e} \alpha \exp(-2(\sigma - 1/\sigma^2)) = \quad (54)$$

$$6.67384 \times 10^{-8} \text{ dyne-cm}^2 - g^{-2}$$

this expression is within 3.6 parts per 100 thousand of the measured value of G:

$6.67408 \times 10^{-8} \text{ dyne-cm}^2 \text{ gm}^{-2}$ . Note that the expression gives proper limiting behavior at the Planck scale, yielding G even as all masses go to  $M_P$ ,  $e^2 \rightarrow \hbar c$  and  $\alpha$  and  $\sigma \rightarrow 1$ .

Therefore, a simple mass model, bridging the lepton-baryon mass system at its lowest energy end-members the electron and proton, that fulfills the expectation of our Gedanken experiment and has proper limiting behavior at both the Planck scale and scale of the fifth dimension, which is the subatomic scale, yields accurate expressions for both the proton mass and the gravitation constant.

A formula similar to Eq. 54 was originally published in approximate form in 1987 and corrected in 1988 [1] and bears some resemblance to the formula published by T'Hooft [16] based on "Instanton" theory that combines Hawking Evaporation with Thermal physics.

### C. Exchange Boson Masses From Quantum Resonant Scattering

The existence of a hidden 5<sup>th</sup> dimension in an otherwise 4 dimensional space time breaks the scale symmetry of the vacuum by inserting a length at which physics must change. Since the 5<sup>th</sup> dimension is independent of the other coordinates, the 5<sup>th</sup> dimension looks like a spherical particle from a distance in any direction, that is, it looks like a particle of a certain size. It is a well-known phenomenon in physics that particles of well-defined sizes in otherwise uniform media support Mie scattering, that is, they support both radial and surface resonances. At the suggestion of Dr. Eric Davis (Private Communication) the consequences of such structural resonances were explored.

Mie scattering would be expected on a hidden dimensional structure in the presence of the ZPF and would give rise to particle quanta. We will also consider the classical particle surface of charged particles as a spherical surface that can support Mie structural resonances. This seems, at first, very unlikely, even bizarre. However, nature appears to do this. The classical surface of a charged particle appears, at first glance, to be a mathematical artifice and not to define a real dynamic entity. However, since this is quantum mechanics, even seemingly unlikely and bizarre events can contribute to observables. This also shows the underlying electromagnetic character of the short range forces.

We will call the particles caused by these quantum Mie scattering events "Mieons." Two fields are available in the ZPF to drive quantum Mie scattering, these are the EM field and Radion field, which must come into being as part of the Kaluza-Klein scheme for having both EM and Gravity and which has the source term  $E^2 - B^2$  [17]. We will identify EM resonances with the factor  $1/\alpha$  and we will identify the Radion resonances with the factor  $\sigma$ . This will give rise to new particles, Mieons, at resonances on the hidden dimension. The EM resonances will be vector resonances around the circumference of the spherical classical surface. The Radion field, being a scalar field, would be expected to produce, at least in lowest order, a simple radial mode inside the spherical classical particle surface. It will also give rise to Mieons on resonances on the classical electrostatic radii of the electron and proton, which will behave like conducting surfaces to first order. The fundamental and lowest order resonances can be expected to be most important as determined by radial and circumferential resonances. The fundamental resonance will be considered as well as a 5-fold resonances because the 5 dimensionality of the entire system for low intensity oscillations.

Since the concept of a quantum resonant path on a classical charged particle surface seems to be but one of many quantum possibilities, we will generalize it to include alternative paths of lower quantum probability, in orders of our coupling strengths,  $\alpha$  and  $1/\sigma$ . Therefore, we express mathematically this concept of Mie resonances, generalized to include virtual paths of reduced probability of order  $\alpha$ , for the EM ZPF, or  $1/\sigma$ , for the Radion field, by the following, for each spin component of the boson:

$$E \ell_{EM} / c = N h \quad (55)$$

$$E \ell_R / c = N \sigma \frac{e^2}{c} \quad (56)$$

Where we have for the path lengths  $\ell$  for EM ZPF excitations, where N and P are integer multiples of  $1/2$ , and express for the Mie quanta agency:

$$\ell_{EM} = (2\pi r_c + \alpha P 2\pi r_c) \quad (57)$$

And likewise for Radion excitations, which are radial:

$$\ell_R = (2r_c + (P/\sigma) 2r_c) \quad (58)$$

Where E is the particle rest energy, c, is the speed of light h is Planck's constant. Rearranging we obtain for the EM ZPF:

$$E = \frac{N h c}{(2\pi r_c + \alpha P 2\pi r_c)} = \frac{N h c}{r_c (1 + \alpha P)} \quad (59)$$

$$E = 2Nm_c c^2 / \alpha(1 + \alpha P) \quad (60)$$

Where  $m_c$  is the particle mass generating the classical radius

$$E\ell / c = N\sigma \frac{e^2}{c} \quad (61)$$

$$\ell = (2\pi r_c + (P / \sigma)2\pi r_c) \quad (62)$$

$$E = 2\sigma Nm_c c^2 / (1 + P / \sigma) \quad (63)$$

The scattering of quanta out of the ZPF by a particle will imprint the quanta with the character of the particle form which it scatters. It must give a spin-state dimensionality of a scalar spin-0 particle off the time-like-scalar nature of the electron-proton at splitting, and a vector spin-1 particle off the space-like proton-quark triplet once it has split. The simplest scatterings will be reactive charge state off the charges of the electron-proton system.




Fig. 6. A quantum Mie scattering caused by a fundamental resonant excitation vector on a classical particle surface




Fig. 7. A quantum Mie scattering caused by a fundamental Radion ZPF resonant scalar excitation of a classical particle shell.

This is the mass of the charged W-boson which has spin 1 reflecting the dimensionality of the proton as having 3 sub-dimensions. These formulas are quite accurate as is seen in Table 1. In both cases the spin of the Mieion is the spin on the classical surface of the

parent particle is plus or minus  $\hbar/2$ . We can take the ratio of the masses of these two bosons:

$$\frac{M_W}{m_{\pi\pm}} = \sigma^3 \alpha = 574.3 \quad (64)$$

We can invert this formula relating the masses of the charged Pions and W boson and use the quantum EM-normalized actual values of their actual masses [18] to derive the value of  $\sigma$ .

$$\sigma = \left( \frac{M_W}{\alpha m_{\pi\pm}} \right)^{1/3} = 42.894 \quad (65)$$

This result is within a part per thousand of the actual value of  $\sigma$ .

TABLE 1. Particle Masses Predicted by the GEM theory for Simple Mie , P=0, Scattering Theory and Observed Masses

Particles	Particle Properties		
	Predicted Mass	Measured Mass	% error
$\pi\pm$	140.05MeV	139.6MeV	0.3%
$W\pm$	80.409GeV	80.398GeV	0.01%
$\eta_c$	3000.6MeV	2985MeV	0.7%

We understand from this that the charged nature of the particles results from the polarization of the vacuum at the classical particle surfaces of the electron and proton. The spin states of the pion and W particles reflect the dimensionality of the electron as a one dimensional particle-yielding a scalar pion, and the three dimensional spin 1 vector character of the W particle is required for it to interact with the three quarks.

We can consider that the path integrals on classical particle surfaces “tumble” in 5 space and to first order all the degrees of freedom are identical. This will allow 5-fold perturbations to develop on the path integral so that virtual paths exist that add or subtract to the effective length of the path, (see Figure 8 and 9) so we will have  $M=5$ .




Fig. 8. A quantum Mie scattering caused by a resonant excitation on a classical particle surface plus a five-fold alternative quantum path.




Fig. 9. A quantum Mie scattering caused by a resonant excitation on a classical particle structure with a 5-fold alternative path also being excited.

We can look thus propose a similar “tumbling in 5 dimensions” process operating in the Radion field except that in case it gives a negative “backflow” or “shortcut” contribution to the path integral.

We return to the path integral model for Mieions generated by the Radion field where  $m_c$  the particle mass generating the classical radius, in this case  $m_e$

$$E = \frac{2\sigma N m_o}{(1 + P/\sigma)} \quad (66)$$

Where we have  $P=-5$   $N=1$  scattering off the proton. As before, the particle must be a spin 1 vector boson because of the “vector” or triune character of the proton. This perturbation features the “backflow”, or “shortcut” negative contribution to the path integral:

$$m_z = \frac{2\sigma m_p}{(1 - 5/\sigma)} = 91.03 GeV/c^2 \quad (67)$$

The perturbed pathway mechanism generally takes charge off of particles because is an averaging over neutral space.

Following this procedure for  $N=1/2$   $M=0$  we obtain for the scalar  $\sigma$  excitation off the uncharged scalar eta-c particle with the uncharged scalar result

$$m_H = \sigma m_{\eta_c} / (1 + 5\alpha) = 124.1 GeV/c^2 \quad (68a)$$

We also obtain EM ZPF excitation off the proton classical surface at spin 0:

$$m_H = m_p / \alpha(1 + 5\alpha) = 124.1 GeV/c^2 \quad (68b)$$

These both give the approximate mass of the Higgs Boson [19] of spin 0 and charge 0. This result (as  $m_H \cong m_p/\alpha$ ) was obtained and presented at the 2012 STAIF Meeting in Albuquerque NM , four months before it was known [4].

So that we have approximately

$$\frac{m_H}{m_{\eta_c}} \cong \sqrt{\frac{m_p}{m_e}} = \sigma = 42.8503 \quad (69)$$

The experimental value is  $m_H/m_{\eta_c} \cong 41.9$ , so agreement is good, within 2%.

Finally, we have for  $N=1/2$  and for  $P=0$  a Radion scattering off the electron, which is a new particle.

$$m_o = \sigma m_e = 21.9 MeV \quad (70)$$

This is the mesoscale particle, which we will term the  $M^*$  (“Morningstar”) particle in honor of the sponsors of this research. It has never been observed, but some evidence for its existence can be found, and will be discussed in the next section of this chapter. We would expect it to be charge-neutral and have spin 0 like the Higgs Boson. It should decay into electron-positron pairs and photons.

The predicted particle masses and those experimentally observed [18] are summarized in Table 2.

TABLE 2. Particle Masses Predicted by the GEM theory and Observed Masses Including the New Predicted  $M^*$  Particle

Particles	Particle Properties		
	Predicted Mass	Measured Mass	% error
$\pi_0$	135.12MeV	134.98MeV	0.1%
Z	91.03GeV	91.19GeV	0.2%
Higgs	124.1GeV	125.1GeV	0.8%
$M^*$	21.98MeV	****	***

III. ELECTROWEAK THEORY AND GEMS  
 CORRESPONDENCE

A complete theory of particle mass creation must include gravity. Accordingly, the Standard Model of Particle Physics must be extended to include gravity. This occurs in the GEMS theory because the Kaluza-Klein schema, in allowing a compact 5<sup>th</sup> dimension to allow EM and gravity fields, must create a scalar field as the element of the metric that controls the 5<sup>th</sup> dimension's measure of itself. This scalar field, called the "Radion" field, that must occur due to a hidden 5<sup>th</sup> dimension, creates both mass and charge in a mathematically similar way as the Higgs Boson.

In Kaluza-Klein, a hidden 5<sup>th</sup> dimension allows gravity and EM fields to coexist, but in GEMS, spacetime is itself electromagnetic, so the existence of a hidden dimension of a certain size creates mass and charge by breaking the scale symmetry of the vacuum, which is itself electromagnetic. GEMS makes the vacuum full of EM fields and "granular", so that if the vacuum is disrupted at a certain or below, particles and the intense EM fields associated with them appear. This in contrast to a classical vacuum without granularity: slice a classical vacuum as finely as one likes, and no EM fields or particles appear. The appearance of the hidden 5<sup>th</sup> dimension creates a scalar field to measure itself, and the fact that the 5<sup>th</sup> dimension has a scale size, creates the subatomic scale where particle EM fields are resolved. The appearance of the 5<sup>th</sup> dimension thus creates particle EM fields and mass associated with them. Thus, the "deployment" or, "compactification" in macroscopic terms, of the 5<sup>th</sup> dimension from the Planck scale created a cosmos full of massive charged particles from the vacuum. In actuality, in the early universe, all dimensions were effectively "compact." The compact nature of the 5<sup>th</sup> dimension only became apparent as the universe expanded, the 5<sup>th</sup> dimension simply deploying to a certain size and then stopping, while everything else expanded.

In GEMS theory, the deployment of the 5<sup>th</sup> dimension from the Planck scale was the trigger of the Big Bang.

This can be seen mathematically by a simple model of a universe filled with a massless quanta, that is a vacuum, with a metric tensor [20]:

$$\gamma_{ab} = \begin{bmatrix} g_{\nu} + \varphi \kappa^2 A_r A_\nu & \varphi \kappa A_r \\ \varphi \kappa A_\nu & \varphi \end{bmatrix} \quad (71)$$

Where  $A$  is the electric 4 potential the indices,  $a$  and  $b$ , run from 1-5,  $\kappa^2=16\pi G/c^4$ ,  $g_{\nu}$  is the familiar 4 metric tensor and  $\varphi$  is a scalar Radion field that

determines the size of the 5<sup>th</sup> dimension, with  $\varphi = 0$  being a primordial 4<sup>th</sup> dimensional universe. We have then a Lagrangian for a set of massless scalar quanta, with  $M_P = (\hbar c/G)^{1/2}$  being the Planck mass, a mass that can arise spontaneously out of the vacuum:

$$L = \frac{\hbar^2}{M_P} \int (\partial_a \phi \partial_b \phi \gamma^{ab}) \sqrt{-\gamma} d^5 x \quad (72)$$

The minimization of this Lagrangian leads to a simple Klein-Gordon wave equation for massless quanta in flat space<sup>6</sup>:

$$\partial_a \partial_b \phi \gamma^{ab} = 0 \quad (73)$$

However, if we allow the fifth dimension to deploy from the Planck Scale and deploy at some large size so that all dependence on the 5<sup>th</sup> dimension becomes the form:

$$\phi = \sum_n \phi_n(x^\nu) \exp(in x^5 / (2\pi r_o)) \quad (74)$$

Where  $n$  is an integer index and  $r_o$  is the size of the hidden dimension. Then the particles in the quantum Lagrangian acquire both mass and charge:

$$L = \frac{\hbar^2}{M_P} \int \left[ \left( \partial_a - \frac{in\varphi}{2\pi r_o} \kappa A_a \right) \phi \right]^2 - \left( \frac{n}{2\pi r_o} \right)^2 \varphi \phi \sqrt{-\gamma} d^5 x \quad (75)$$

Where the electric charge is identified as:

$$q = \frac{\varphi n \kappa}{2\pi r_o} \quad (76)$$

$$m = \frac{\hbar \sqrt{\varphi}}{2\pi r_o c^2} \quad (77a)$$

So that for field free space, we have:

$$\partial_a \partial_b \phi \gamma^{ab} + \left( \frac{m c}{\hbar} \right)^2 \varphi \phi = 0 \quad (77b)$$

So that the compactification of a 5<sup>th</sup> dimension allows both charged and massive particles to appear from a previously vacuum filled universe and creates a Kaluza-Klein Radion scalar field to do this. Therefore, a Kaluza-Klein model creates mass and gravity in the same formalism and thus automatically solves the

Luhon conjecture “ One cannot create mass without creating gravity”. This Radion-Higgs correspondence has also been proposed independently by Wesson [21]

This appearance of a hidden 5<sup>th</sup> dimension of a specific size corresponds to SSB “Spontaneous Symmetry Breaking” in Standard Electroweak theory[22]. The existence of massive particles can thus be explained by scalar Higgs field and SSB but also by the appearance of a 5<sup>th</sup> dimension and its compactification

In Standard Model Electro-Weak theory the Universe begins with a SSB event that creates a new vacuum state with a Higgs mass and a consequent Vacuum Expectation Value that creates mass in other particles . An SU(2) symmetry exists which rotates a B and W<sub>3</sub> set of fields into a massless photon  $\gamma$ , and Z boson.

$$\begin{pmatrix} \gamma \\ Z \end{pmatrix} = \begin{pmatrix} \cos\theta_w & \sin\theta_w \\ -\sin\theta_w & \cos\theta_w \end{pmatrix} \begin{pmatrix} B \\ W_3 \end{pmatrix} \quad (78)$$

where  $\theta_w$  is the “Weak Mixing Angle”

The value of  $\theta_w$  is set by the ratio of the masses of the charged W and neutral Z bosons.

$$\cos\theta_w = \frac{M_w}{M_z} = 0.88165 \quad (79)$$

These masses are not given explicitly in the Electro-Weak theory but are given by the GEMS theory so we can write, from Eq. 79:

$$\cos\theta_w = (1 - \frac{5}{\sigma}) \quad (80)$$

$$\sin\theta_w = \frac{5}{\sigma} \sqrt{2\sigma/5 - 1} \quad (81)$$

Note that  $\cos\theta_w = 1$  and  $\sin\theta_w = 0$  when  $\sigma = 5$ , near the Planck Scale. We have then from the observed masses of the M and Z bosons a value for  $\sigma$  in our present universe:

$$\sigma = \frac{5}{1 - \frac{M_w}{M_z}} = 42.2489 \quad (82)$$

This result is within 1.4% of the actual value of  $\sigma$ . Thus the GEMS parameter already is operating in the Standard Model.

It is apparent from this that the energy of EM-Weak Force Unification (EWU) occurs quite close to the Planck Energy, at  $\sigma = 5$ . Based on Eq. 44, this gives an energy scale  $E_{EWU}$  of approximately  $10^{17}$  GeV:

$$\ln \left| \frac{E_{EWU}}{E_p} \right| = 4.75 \quad (83)$$

where  $E_p$  is the Planck energy of  $1.22 \times 10^{19}$  GeV. However, Strong Force-EM-Weak Force unification does not occur in GEMS until  $\sigma=1$ , that is, at the Planck Energy, explaining the proton’s stability.

TABLE 3. Correspondence between the GEMS and Electro-Weak Unification theories.

Electro-Weak- GEMS Correspondence		
Electro-Weak Theory	GEMS Unification	comments
Spontaneous Symmetry Breaking	5 <sup>th</sup> Dimension Deployment	Big Bang
Higgs field	Kaluza-Klein Radion field comes into existence with 5 <sup>th</sup> dimension	gravity, charge, and mass are born together
SU(2) symmetry	SO(3) symmetry	Isomorphic groups
$\theta_w$ rotation evolution	$\sigma$ evolution	$\sigma \rightarrow 1$ at Planck Scale
Unification energy $\sim 10^{16}$ GeV	Electro Weak unification at $\sim 10^{17}$ GeV	Strong-Weak Unification at $1.22 \times 10^{19}$ GeV (Planck scale)

Unexpectedly, the GEM theory created a doorway to understanding with two short-range forces of nature the Weak and Strong nuclear forces, because in unifying gravity and EM in a geometric theory, it produced a geometric scale regime for subatomic particles and the regime for their interactions. The quantum particles which create the short range forces are thus scatterings out of the full spectrum of the ZPF by these resonant structures. The fact that the scattering structures are EM classical radii shows the underlying electromagnetic character of the short range forces. The GEM theory produced the picture of EM forces not only between charged objects but also

between uncharged structures that can be extended to include short-range nuclear forces.

#### IV. SUMMARY AND DISCUSSION

Under the conjecture of Dr. Alfred Luhen, (Private Communication) 'one cannot create mass without creating gravity.' The Higgs Boson thus cannot exist and generate mass outside the context of General Relativity. Accordingly, the simplest way for this to occur in the GEM context is that the Higgs scalar field occurs in Kaluza-Klien theory as the Radion scalar field, so that both gravity, EM-mass energy as gravity source term, and particle charges and masses are born together with the Radion field. The concept of the Higgs Boson as the creator of mass in the GEMS theory is obvious because of the relationship  $m_{\text{Higgs}} \cong m_p/\alpha$  so that the Compton radius of the Higgs Boson is the EM interaction length of the proton:

$$\hat{\lambda}_{\text{Higgs}} \cong \frac{e^2}{m_p c^2} \quad (84)$$

Thus, the known source of mass in the universe, the proton, is in EM resonance with the Higgs Boson, that is, the proton EM self-interaction time is the Compton oscillation time of the Higgs. The Higgs boson can thus be viewed as the most general excitation, by both EM and Radion fields, of a structural resonance of the hidden Kaluza-Klien 5<sup>th</sup> dimension, and thus part of the mechanism in the vacuum that gives rise to separate EM and gravity fields and also a cosmos dominated by hydrogen.

The splitting of the four forces mirrors the splitting of the proton-electron pair in the GEMS theory, with the deployment of the hidden fifth dimension. The splitting of the Strong Force from the Electro-Weak Force occurs early, at the Planck scale, with the splitting of the photons into the gluon octet. The splitting of EM from the Weak Force at  $\sim 1/100$  of this energy.

The GEMS theory has deep correspondence to the Standard Model, especially the Electro-Weak Unification theory, because of its mass-creating Higgs process and its unifying character. However, whereas the Electro-Weak theory cannot predict masses directly, but only their ratios, the GEMS predicts particle masses. The important ratio of the W boson mass to the Z boson mass, in particular, is derived accurately and illustrates the presence of the GEMS mass parameter  $\sigma = 42.8503$  in the Standard Model even through its presence is unrecognized. The central importance of the EM classical radii in this unification theory suggests the underlying EM character of the forces. The model gives the correct masses, spins and charges reflecting the dimensionality of the electron or proton they scatter off of. The

The result is a rudimentary "Bohr Model" of field unification which gives G, the mass of the proton, and the masses of the pions and W and Z exchange bosons of the Strong and Weak force. It also gives an accurate estimate of the Higgs Boson mass. It also predicts new particles at 21 and 22 MeV and other phenomena, particularly that hydrogen and radiation can appear occasionally from the vacuum, particularly at Black Hole mergers [5]. This theory suggests that manipulation of Gravity, Strong, and Weak Forces by Electromagnetism may be possible. It is hoped this work can form the basis for future advances in understanding and engineering.

#### ACKNOWLEDGMENT

The author wishes to thank Morgan Boardman and Paul Murad of Morningstar Applied Physics and Eric Rice of Orbital Technologies Corporation and Jess Sponable of DARPA for their support and encouragement of this research as well as Abe Meghed for many useful comments, and finally thanks to my industrious cousin Axel for his good example. The author is also very grateful to JMESS for this opportunity to codify the GEM theory in its present, early, state. *Laus Deo*

#### REFERENCES

- [1] Brandenburg J.E. (1988) "Progress on the GEM ( Gravity-Electro-Magnetism) Theory" APS Bull., 32, 1, p. 32.
- [2] Brandenburg J.E. (1992) "Unification of Gravity and Electromagnetism in the Plasma Universe" IEEE Transactions on Plasma Science, Vol 20, 6, p944.
- [3] Brandenburg, J. E., "A Model Cosmology Based on Gravity-Electromagnetism Unification", Astrophysics and Space Science, **227**,133-145, (1995).
- [4] Brandenburg, J.E., "An Extension of the GEM Unification Theory to Include Strong and Weak Nuclear Forces and an Estimate of the Higgs Boson Mass STAIF II Conference Albuquerque NM March 2012. Published in Journal of Space Exploration, Vol 1, Issue1 (2012)
- [5] Brandenburg, J.E. "The GEM Unification Theory: Extending The Standard Model to Include Gravitation" Lambert Academic Publishing (2016).
- [6] ] Brandenburg, J. E., "The GEM (Gravity-EM) Theory:the Unification of the Strong, EM ,Weak ,and Gravity Forces of Nature" Journal of Multidisciplinary Engineering Science Studies (JMESS) ISSN: 2458-925X Vol. 2 Issue 7, July – 2016



- [7] J. L. Feng, B. Fornal, I. Galon, S. Gardner, J. Smolinsky, T. M. P. Tait, and P. Tanedo, "Evidence for a Protophobic Fifth Force from 8Be Nuclear Transitions," arXiv:1604.07411 [hep-ph].
- [8] Klein, Oskar: "Quantum Theory and Five-Dimensional Relativity Zeitschrift fur Physik, 37, 895, (1926).
- [9] Sakharov A.D.:(1968) "Vacuum quantum fluctuations in curved space and the theory of gravitation" Sov. Phys. Doklady 12,p.1040-104.
- [10] "Introduction to Plasma Physics" , Francis Chen,(1974) p34.
- [11] Zel'dovich, Y.B., JETP letters **6**, 316 (1967); "The Cosmological Constant and Elementary Particles" , Soviet Physics Uspekhi **11**, 381 (1968)
- [12] Puthoff, H.E., "Gravity as a zero-point fluctuation force", *Phys. Rev. A* **39**, 2333, (1989).
- [13] Brandenburg, John, Derivation of the Schwartzchild Metric From the "Self Censorship" of the ZPF (Zero Point Energy) in the GEM Theory, Fall 2012 Meeting of the APS Ohio-Region Section, October 5-6, 2012, abstract #EC.008
- [14] Marmier P. and Sheldon E., *Physics of Nuclei and Particles* (Academic Press, New York, 1969), vol. 1.
- [15] Hawking, S. W. (1974). "Black hole explosions?". *Nature* **248** (5443)
- [16] t'Hooft G., "A physical interpretation of gravitational instantons." Nucl. Phys. **B315** (1989) 2, 517-527.
- [17] Williams L.L. "Field Equations and Lagrangian for the Kaluza Metric Evaluated with Tensor Algebra Software" Journal of Gravity, Volume 2015, Article ID 901870
- [18] K.A. Olive et al. (Particle Data Group), Chin. Phys. C38, 090001 (2014)
- [19] Evidence for the direct decay of the 125 GeV Higgs boson to fermions, The CMS Collaboration, *Nature Physics*, **10**, 557-560 (2014)
- [20] Mann, R.B. (1984) "Calculation of Coupling Constants In Kaluza-Klein Theories," from "An Introduction to Kaluza-Klein Theories" Lee H. C., Editor, World Scientific Publishing Co. Ltd. p.233
- [21] Wesson, P.S. (2010) "The Scalar Field Of 5D Gravity And The Higgs Field Of 4D Particle Physics:APossibleConnection,"arXiv:1003.2476v 1 [gr-qc]
- [22] Odintsov, S. D., (1988) "Compactification and spontaneous symmetry breaking in the  $\lambda\phi^4$  theory with Kaluza-Klein background" Soviet Physics Journal, Vol. 31, Issue 9, pp 695-699.