

Statistical Converges of Series and Statistical Pettis Integration

¹Anita Caushi

Department of Mathematical Engineering
 Polytechnic University of Tirana
 Tirana, Albania
 anita_caushi@yahoo.com

²Ervenila Musta

Department of Mathematical Engineering
 Polytechnic University of Tirana
 Tirana, Albania
 ervimst@yahoo.com

Abstract—Pettis and Dunford's integrals are the most important concepts concerning the modern theory of probabilities. We extend this to a statistical form. In this paper, we prove the countable additive of the statistical Pettis integral. For, this we need some properties of the unconditional statistical convergence of series in Banach spaces. We give an example where we show' if a series is statistically unconditionally convergent then it is weakly absolutely statistically convergent.

Keywords—*st-convergence; st-measurability st-weakly measurability; st-integrability*

I. INTRODUCTION

IT IS KNOWN THAT THE IDEA OF STATISTICAL CONVERGENCE WAS GIVEN BY ZIGMUND [12]. THE CONCEPT OF STATISTICAL CONVERGENCE WAS FORMALIZED BY STEINHAUS[11] AND FAST [6]. SOME YEARS LATER, THE CONCEPT WAS REINTRODUCED BY SCHOENBERG [9]. STATISTICAL CONVERGENCE HAS BECOME AN ACTIVE AREA OF RESEARCH IN RECENT YEARS. IN THIS PRESENTATION, WE FOLLOW CONCEPTS INTRODUCED BY FRIDY[8] ABOUT THE CONVERGENCE OF SEQUENCES AND THE CONCEPT OF SCHOENBERG ABOUT INTEGRATION THE BASIC CONCEPT IS THE STATISTICALLY CAUCHY CONVERGENCE OF FRIDY[7].

II. PRELIMINAIRES

Let be A_n a subset of an ordered natural set N . It said

to have a density $\delta(A)$ if $\delta(A) = \lim_{n \rightarrow \infty} \frac{|A_n|}{n}$, where

$A_n = \{k < n : k \in A\}$ and with $|A|$ denotes the cardinality of the set A . It is clear that the finite sets have the density zero and $\delta(A') = 1 - \delta(A)$ if $A' = N - A$. If a property $P(k) = \{k : k \in A\}$ holds for all $k \in A$ with $\delta(A) = 1$, we say that property P holds for almost all k that is a.a.k. The vectorial sequence x is statistically convergent to the vector(element) L if for each $\varepsilon > 0$

$$\lim_{n \rightarrow \infty} \frac{1}{n} |\{k \leq n : \|x_k - L\| \geq \varepsilon\}| = 0$$

or, $\|x_k - L\| < \varepsilon$ a.a.k

Definition 1. A series of elements $x_k \in X, k \in N$ of a Banach space X is said to be statistical convergent if

the sequence of its partial sums $S_n = \sum_{k=1}^n x_k$ statistical convergent in the norm of the space X .

Definition 2 . The series $\sum_{k=1}^{\infty} x_k, x_k \in X, k \in N$ is absolutely statistically convergent if $\sum_K \|x_k\| < \infty, k \in K \in N, \delta(K) = 1$.

Proposition 1. If $\sum_{k=1}^{\infty} x_k$ converges statistically absolutely then $\sum_{k=1}^{\infty} x_k$ is statistical convergent.

Definition 3. A series $\sum_{k=1}^{\infty} x_k$ of elements $x_k \in X, k \in N$ of a Banach space X is said to be statistical unconditionally convergent if is statistically convergent for every rearrangement of its terms, i.e. if the series $\sum_K x_{p(k)}, k \in K \in N, \delta(K) = 1$ converges whenever P is a one-to-one mapping of N onto N .

Theorem. For a series $\sum_{k=1}^{\infty} x_k$ of elements $x_k \in X, k \in N$ of a Banach space X the following conditions are equivalent:

- (a) the series converges statistically unconditionally
- (b) all series of the form $x_{n_1} + x_{n_2} + x_{n_3} + \dots$ where $n_1 < n_2 < n_3 < \dots$ statistical convergent,
- (c) for every bounded sequence $(a)_i, (a)_i \in R$ the series $\sum_K a_k x_k$ statistical convergent to some element of X .

Proposition 2.

If $\sum_{k=1}^{\infty} x_k$ of elements $x_k \in R, k \in N$ is statistical unconditionally convergent if and only if is statistically absolutely convergent.

Definition 4.

A sequence $x_k \in X, n \in N$ statistically weakly converges to $x \in X$ if for every $x^* \in X^*$

$$\lim_K x^*(x_k) = x^*(x)$$

Definition 5 .

A series $\sum_{k=1}^{\infty} x_k, x_k \in X, k \in N$ statistical weakly convergent to a sum $s \in X$ if for every $x^* \in X^*$ the limit

$$\lim_K x^*(\sum_K x_k) = \lim_K x^*(\sum_K x_k) = x^*(s)$$

Theorem. (Orlicz, Pettis)[10]

Let $\sum_{k=1}^{\infty} x_k, x_k \in X, k \in N$ a series in a Banach space X .

If for each set $A \subset N$ there is $x_A \in X$ such that for each $x^* \in X^*$ we have $\sum_{k \in A} x^*(x_k) = x^*(x_A)$ then the series $\sum_{k=1}^{\infty} x_k$ is unconditionally convergent.

Definition 6 :

A series $\sum_{k=1}^{\infty} x_k$ is called statistical weakly absolutely convergent if $\sum_K |x^*(x_k)| < \infty$

Proposition 3 .

If $\sum_{k=1}^{\infty} x_k$ is statistically unconditionally convergent then is statistical weakly absolutely convergent.

Example.

For $k \in N$ denote by

$$e_k = \begin{cases} (0, \dots, 0, k, 0 \dots) & k \text{ prime} \\ (0, \dots, 0, 1, 0, 0 \dots) & \text{others} \end{cases}$$

e_k is the element of c_0 with 1 on the k -th position in the sequence.

Since we have

$$\sum_{k=1}^n e_k = \begin{cases} (1, 2, 3, 1 \dots n, 0, 0 \dots), & k \text{ prime} \\ (1, 1, 1 \dots 1, 0, 0 \dots), & \text{others} \end{cases}$$

For $N \in N$, we can see immediately that the series $\sum_{k=1}^{\infty} e_k$ does not converge in c_0 (in the norm)

and the serie $\sum_K e_k$ has the same nature.

Assume that $x^* = c_0^* = l_1$, i.e. $x^* = (\alpha_1, \alpha_2, \dots, \alpha_k, \dots)$

, $\alpha_k \in R, k \in N$

st- $\sum_{k=1}^{\infty} |\alpha_k| = \|x^*\|_{c_0^*} < +\infty$ or $\sum_K |\alpha_k| < +\infty$.

Then $x^*(e_k) = \alpha_k$ for k not prime

$$\sum_K |x^*(e_k)| = \sum_K |\alpha_k| < +\infty$$

i.e the series $\sum_{k=1}^{\infty} e_k$ is stastically weakly convergent .

Since $x^*(\sum_K e_k) = \sum_K \alpha_k$ for $k \in K$ we can see that this sequence converges to $\sum_K \alpha_k = x^*(y)$

, where $y = (0, 0, 1, 0, 1, 0, 1, 1 \dots)$, is a sequence which does not belong to c_0 . This means that if the series $\sum_{k=1}^{\infty} e_k$ were statistically weakly convergent then its weak sum would be y but c_0 contains such element.

Definition 7 .A weakly measurable $f : S \rightarrow X$ with $x^*(f)$ Lebesgue integrable for every $x^* \in X^*$ is Statistical Pettis integrable if for every measurable $E \subset S$ there is an element $x_E \in X$ that satisfies

$$x^*(x_E) = \int_E x^*(f)$$

for every $x^* \in X^*$.

Theorem .

If $f : S \rightarrow X$ is st- Pettis integrable define for a measurable set $E \subset S$ the function

$\vartheta(E) = Ps - \int_E f d\mu \in X$ (the indefinite st-Pettis integral).

The function ϑ is countably additive.

Proof:

Assume that $E_n \subset S, n \in N$ are measurable sets, $E_n \cap E_m = \emptyset, n \neq m$. Then

$$x^*(\vartheta(\bigcup_{n=1}^{\infty} E_n)) = x^*(Ps$$

$$- \int_{\bigcup_{n=1}^{\infty} E_n} f d\mu = Bs - \int_{\bigcup_{n=1}^{\infty} E_n} x^*(f) d\mu$$

$$= \sum_{n=1}^{\infty} (Bs - \int_{E_n} x^*(f) d\mu) = \sum_{n=1}^{\infty} x^*(\vartheta(E_n))$$

For every $x^* \in X^*$. This means that \cup is weakly countably additive, i.e. the series of real numbers $\sum_{n=1}^{\infty} x^*(\vartheta(E_n))$ is convergent for every $x^* \in X^*$. Hence it is also unconditionally convergent this means that it is also weakly subseries convergent. The Orlicz-Pettis theorem [10] yields that the series $\sum_{n=1}^{\infty} \vartheta(E_n)$ is statistically unconditionally convergent and henceforth convergent in norm.

While $\sum_{n=1}^{\infty} \vartheta(E_n) = \vartheta(\bigcup_{n=1}^{\infty} E_n)$

The theorem is proved.

Theorem. Let $f : S \rightarrow X$ be statistical measurable of the form

$$f = g + \sum_K x_n \chi_{E_n} \quad \delta(K) = 1 \quad (1)$$

where $g : S \rightarrow X$ is measurable and bounded, E_n are pairwise disjoint measurable subsets of $S, x_n \in X, n \in N$ and $E_n, n \in N$ can be chosen such that the series $\sum_{n=1}^{\infty} x_n \mu(E_n)$ statistical converges unconditionally in X , and in this case we have

$$Ps - \int_E f d\mu = Ps - \int_E g d\mu + \sum_K x_n \mu(E \cap E_n) \quad (2)$$

for every measurable $E \subset S$ and $\delta(K) = 1$

Proof. Assume that f is statistical Pettis integratin of the form (1). Since g is statistical bounded we have $g \in B_S \subset P_S$ by roposition 3.9 [10] and therefore also

$$h = f - g = \sum_K x_n \chi_{E_n} \in P_S$$

If $E \subset S$ is measurable then, because the indefinite statistical Pettis integral is countably additive by Theorem 3.24, [10], we have

$$\int_E h d\mu = \sum_K \int_{E \cap E_n} h d\mu = \sum_K x_n \mu(E \cap E_n) \quad , \delta(K) = 1$$

Taking any rearrangement of the series $\sum_K x_n \chi_{E_n}$, we obtain the same function h , i.e.

$$h = \sum_K x_n \chi_{E_n} = \sum_{k=1}^{\infty} x_{\pi(n)} \chi_{E(\pi(n))}$$

for any one-to-one map π of K onto N and of course

$$\int_E h d\mu = \sum_{n=1}^{\infty} \int_{E \cap E_n} h d\mu = \sum_{k=1}^{\infty} x_{\pi(n)} \mu(E \cap E_n) \in X$$

Hence the series $\sum_{n=1}^{\infty} x_n \mu(E_n)$ is unconditionally convergent.

To show the converse let us mention that the function $g : S \rightarrow X$ being bounded ($\|g(t)\|_X \leq M$ for almost all $t \in S$) is statistical Bochner integrable and therefore $g \in P_S$ by Proposition 3. [10] Now it suffices to show that $h = \sum_K x_n \chi_{E_n}$, is statistical Pettis integrable

provided the series $\sum_{n=1}^{\infty} x_n \mu(E_n)$ is unconditionally convergent in X .

Without loss of generality it can be assumed for simplicity that $\mu(E_n) > 0, n \in N$. Assume that $E \subset S$ is measurable.

Then the series

$$\sum_K x_n \mu(E \cap E_n) = \sum_K x_n \mu(E \cap E_n) \frac{\mu(E \cap E_n)}{\mu(E_n)}$$

is unconditionally convergent in X because $\frac{\mu(E \cap E_n)}{\mu(E_n)} \leq 1$, for all $n \in N$ (Theorem 3.21, [10]).

If $x^* \in X^*$ then $\sum_K x_n^* \mu(E \cap E_n)$ converges unconditionally in R and therefore by proposition 3.22 [4] we have

$$\int_E |x^*(h)| d\mu = \sum_K x_n^* \mu(E \cap E_n) = x^*(\sum_K x_n \mu(E \cap E_n))$$

And $h \in P_S$. While

$$P_S - \int_E h d\mu = \sum_K x_n \mu(E \cap E_n)$$

This yields $f = g + h \in P_S$ and also the equality (2)

Theorem 2.3.4. Suppose that X does not contain subspaces isomorphic to C_0 and let $f: S \rightarrow X$ be Statistical Dunford integrable. If f is statistical measurable, then f is Statistical Pettis integrable on S .

Proof. Since f is measurable, we have the relation

$$f = g + \sum_K x_n \chi_{E_n} \quad \delta(K) = 1$$

where $g: S \rightarrow X$ is measurable and bounded, E_n are pairwise disjoint measurable subsets of S , $x_n \in X, n \in N$. Since the interval S is compact, the function $g: S \rightarrow X$ is Bochner integrable and by Proposition 2.3.1, [10] also Pettis integrable. The Statistical Dunford integrability $\sum_K x_n \chi_{E_n}$ yields the Statistical Dunford integrability (Lebesgue integrability) of $x^*(\sum_K x_n \chi_{E_n})$ for every $x^* \in X^*$ and we have also $x^*(\sum_K x_n \chi_{E_n}) = \sum_K x^*(x_n \chi_{E_n})$ because the sets E_n are pairwise disjoint.

Therefore we have

$$\sum_K x^* |x_n \mu(E_n)| < +\infty \text{ for every } x^* \in X^*.$$

This implies that the series $\sum_K x_n \mu(E_n)$ statistical weakly absolutely converges.

Since X does not contain subspaces isomorphic to C_0 , by the Bessaga-Pelczyrski Theorem B.22 [10] presented in Appendix B, the series $\sum_K x_n \mu(E_n)$ converges unconditionally in X . Hence, $\sum_K x_n \chi_{E_n}$ is Pettis integrable by Proposition 2.3.3 [10] and we have $f \in P$.

Theorem .Suppose that X does not contain subspaces isomorphic to C_0 and let $f: S \rightarrow X$

Statistical Dunford integrable. If $Ds - \int_E f d\mu \in X$ for every interval $J \subset S$ then f is Statistical Pettis integrable on S .

Proof. First of all we have the following statement
 If $J_k \subset S, k \in N$ is a sequence of non-overlapping intervals then $Ds - \int_{J_k} f d\mu \in X$
 Indeed, we have

$$B_S - \int_{\cup J_k} x^*(f) d\mu = \sum_K (B_S - \int_{J_k} x^*(f) d\mu) = \sum_K \left| x^*(D_S - \int_{J_k} f d\mu) \right| < +\infty$$

for every $x^* \in X^*$. Hence the series $\sum_K Ds - \int_{J_k} f d\mu$ statistical weakly also lutely converges.

Since X does not contain subspaces isomorphic to C_0 the Bessaga-Pelczyrski Theorem [10] implies that the series $\sum_K Ds - \int_{J_k} f d\mu$ statistically unconditionally converges to a certain element $x_{\cup J_k} \in X$

and $Ds - \int_{\cup J_k} f d\mu = x_{\cup J_k} \in X$, By Theorem (1.11), [10] every open set in $R^n, m \geq 1$ can be written as a countable union of non-overlapping (closed) intervals and therefore by the statement above we obtain that $Ds - \int_G f d\mu \in X$ for every open set $G \subset S$. If $F \subset S$ is closed then $S \setminus F$ is open (in S) and

$$D_S - \int_F f d\mu = (D_S - \int_S f d\mu) - (D_S - \int_{S \setminus F} f d\mu) \in X$$

Note that if $Z \subset S$ is such that $\mu(Z) = 0$ then $Ds - \int_Z f d\mu = 0 \in X$. Let now $E \subset S$ be an arbitrary measurable set. Then by Theorem (3.28) [10] we have $E = H \cup Z$ where $\mu(Z) = 0$ and H is of type F_σ i.e where $H = \cup_k H_k, k \in N$ are closed. Define $L_n = \cup_{k=1}^n H_k$. The sets $H_k \subset S, k \in N$ are closed and $L_n \subset L_{n+1}, n \in N$. Set $L_0 = \emptyset, K_n = L_n \setminus L_{n+1}, n \in N$. Then $K_n \cap K_l = \emptyset$ per $n \neq l$ and $H = \cup_{n=1}^{\infty} K_n$.
 Note that

$$D_S - \int_{K_n} f d\mu = D_S - \int_{L_n \setminus L_{n+1}} f d\mu = \left(D_S - \int_{L_n} f d\mu \right) - \left(D_S - \int_{L_{n+1}} f d\mu \right) \in X$$

Further

$$\begin{aligned} \int_H x^*(f) d\mu &= \int_{\bigcup k_n} x^*(f) d\mu \\ &= \sum_n \int_{k_n} x^*(f) d\mu \\ &= \sum_n x^* \left(Ds - \int_{k_n} f d\mu \right) d\mu \end{aligned}$$

And

$$\begin{aligned} \sum_n \left| x^* \left(Ds - \int_{k_n} f d\mu \right) d\mu \right| &= \sum_n \left| Bs - \int_{k_n} x^*(f) d\mu \right| \leq \\ \sum_n Bs - \int_{k_n} |x^*(f)| d\mu &= \int_{\bigcup k_n} |x^*(f)| d\mu \\ = \int_H |x^*(f)| d\mu &< +\infty \end{aligned}$$

for every $x^* \in X^*$.

Similarly as above the Bessaga-Pelczynski Theorem B.22, [10] implies that the series $Ds - (\int_{k_n} f d\mu)$ statistical unconditionally converges to a certain element $x_H \in X$ and

$$Ds - \left(\int_H f d\mu \right) = x_H \in X$$

Hence

$$Ds - \int_E f d\mu = Ds - \int_H f d\mu + Ds - \int_Z f d\mu \in X$$

References

- [1] Bhardwaj V. Balai., "On Weakly Statistical Convergence", international Journal of Mathematics and Mathematical Sciences, Volume 2007, Article D 38530, 9 pages
- [2] Çakalli H., A study on statistical convergence. Functional analysis, approximation and computation 1:2(2009),19-24
- [3] Caushi, A., Tato, A., A statistical integral of Bohner type on Banach space, HikariLtd Appl. Math. Sci., Vol. 6, 2012, no. 137-140, 6857-6870.
- [4]Caushi ,A. ,Tato.A., A statistical integral of Pettis type on Banach space ,Journal of Advances in Mathematics.Vol 3, No 2. 2013
- [5] Connor J., Ganichev M., and Kadets V., "A characterization of Banach spaces with separable duals via weakly statistical convergence," Journal of Mathematical Analysis and Applications, vol. 244, no. 1, pp. 251–261, 1989.
- [6] Fast H., "Sur la convergence statistique," Colloquium Mathematicum, vol. 2, pp. 241–244, 1951.
- [7] Fridy J. A., "On statistical convergence," Analysis, vol. 5, no. 4, pp. 301–313, 1985.
- [8] Fridy J. A., "Statistical limit points," Proceedings of the American Mathematical Society, vol. 118, no. 4, pp. 1187–1192, 1993.

[9] Schoenberg J., "The integrability of certain functions and related summability methods," The American Mathematical Monthly, vol. 66, no. 5, pp. 361–375, 1959.

[10] Schëabik S.,Guoju Y.Topics in Banach space integration, Series in Analysis vol. 10. world Scientific Publishing Co. Singapore 2005.

[11] Steinhaus H., "Sur la convergence ordinaire et la convergence asymptotique," Colloquium Mathematicum, vol. 2, pp. 73–74, 1951.

[12] Zygmund A., Trigonometric Series, Cambridge University Press, Cambridge, UK, 1979